



Review

The metabolic basis of *Candida albicans* morphogenesis and quorum sensingTing-Li Han^a, Richard D. Cannon^b, Silas G. Villas-Bôas^{a,*}^a Centre for Microbial Innovation, School of Biological Sciences, The University of Auckland, 3sA Symonds Street, Auckland 1142, New Zealand^b Department of Oral Sciences, The University of Otago, Dunedin, New Zealand

ARTICLE INFO

Article history:

Received 23 July 2009

Accepted 5 April 2011

Available online 12 April 2011

Keywords:

Candida albicans

Filamentous growth

Central carbon metabolism

Amino acid biosynthesis

Sterol biosynthesis

Farnesol

Fungi

Dimorphism

Amino acids

Biosynthesis

Systems biology

Functional genomics

Transcriptomics

Proteomics

Metabolomics

ABSTRACT

Candida albicans is a polymorphic fungus that has the ability to rapidly switch between yeast and filamentous forms. The morphological transition appears to be a critical virulence factor of this fungus. Recent studies have elucidated the signal transduction pathways and quorum sensing molecules that affect the morphological transition of *C. albicans*. The metabolic mechanisms that recognize, and respond to, such signaling molecules and promote the morphological changes at a system level, however, remain unknown. Here we review the metabolic basis of *C. albicans* morphogenesis and we discuss the role of primary metabolic pathways and quorum sensing molecules in the morphogenetic process. We have reconstructed, *in silico*, the central carbon metabolism and sterol biosynthesis of *C. albicans* based on its genome sequence, highlighting the metabolic pathways associated with the dimorphic transition and virulence as well as pathways involved in the biosynthesis of important quorum sensing molecules.

© 2011 Elsevier Inc. All rights reserved.

Contents

1. Introduction	748
2. Signaling pathways and transcriptional regulators associated with <i>C. albicans</i> morphogenesis	749
3. Quorum sensing molecules and <i>C. albicans</i> morphogenesis	752
4. <i>In silico</i> reconstruction of the central carbon metabolism of <i>C. albicans</i> and the biosynthesis of quorum sensing molecules	754
4.1. Glycolysis	754
4.2. Pentose phosphate pathway	754
4.3. Pyruvate dehydrogenase	754
4.4. TCA cycle	754
4.5. Glyoxylate shunt	754
4.6. Amino acid biosynthesis	756
4.6.1. Glutamate biosynthesis	756
4.6.2. Aromatic amino acid and quorum sensing molecule biosynthesis	756
4.6.3. Regulation of global amino acid biosynthesis	756
5. Central carbon metabolism, virulence, and morphogenesis of <i>C. albicans</i>	756
6. The effect of metabolic genes on the utilization of carbon sources	757
7. <i>In silico</i> reconstruction of the sterol biosynthetic pathway and farnesol biosynthesis in <i>C. albicans</i>	757
8. The integration of quorum sensing and central carbon metabolism in <i>C. albicans</i>	757
9. Final remarks	758
References	759

* Corresponding author. Fax: +64 9 373 7416.

E-mail addresses: richard.cannon@otago.ac.nz (R.D. Cannon), s.villas-boas@auckland.ac.nz (S.G. Villas-Bôas).

1. Introduction

Candida albicans is a human commensal fungus that can be isolated from approximately 70% of the healthy population (Mavor et al., 2005). In the majority of cases *C. albicans* is harmless, however, if the person is immunocompromised, it can be an opportunistic pathogen (Wingard et al., 1979). *C. albicans* is the fourth leading cause of nosocomial bloodstream infections (Pfaller and Diekema, 2007), with an attributable mortality of 37–44% in severely immunocompromised patients (Wisplinghoff et al., 2004; Leroy et al., 2009; Moran et al., 2010).

C. albicans and other polymorphic fungi (e.g. *Histoplasma capsulatum*) have the remarkable ability to grow in several distinct morphological forms: yeast, hyphae, and pseudohyphae, according to environmental conditions (Bastidas and Heitman, 2009; Sudbery et al., 2004). The true hyphae and pseudohyphae (chains of elliptical cells with constrictions at the septa) are often referred to as filamentous forms (Odds, 1988). The ability to switch rapidly from yeast-to-filamentous growth or vice versa in response to certain environmental cues is considered to be a critical virulence factor for these fungi (Lo et al., 1997; Mitchell, 1998; Brown and Gow, 1999; Gow et al., 2002; Rooney and Klein, 2002; Nemecek et al., 2006). In *C. albicans*, each morphology is believed to confer discrete advantages in the course of infection. The yeast form is important for dissemination through the bloodstream (Bendel et al., 2003; Saville et al., 2003), and adheres to endothelial surfaces (Grubb et al., 2009). The filamentous forms, on the other hand, are more adapted for invasion through the host epithelial tissue (Rooney and Klein, 2002), and also have a higher resistance to phagocytosis due to

their morphology. Indeed, an engulfed *C. albicans* yeast cell can destroy a macrophage if filamentous growth is triggered after phagocytosis (Arai et al., 1977; Lorenz et al., 2004), and filamentous forms have a higher resistance to neutrophil killing (Smail et al., 1992; Fradin et al., 2005). Moreover, experimental studies support the hypothesis that the morphological transition is an essential virulence factor for *C. albicans*. For instance, a reduced mortality rate has been reported in animal infection models for mice inoculated with *C. albicans* mutants unable to undergo the yeast-to-filamentous transition (Lo et al., 1997; Gale et al., 1998). In addition, the induction of hyphal gene expression promoted virulence in a mouse model of systemic candidiasis (Carlisle et al., 2009).

The morphogenesis of *C. albicans* is predominately determined by environmental signaling. The yeast-to-filamentous transition can be triggered by serum, proline, N-acetylglucosamine, different carbon sources, and other cues as summarized in Table 1. However, there is a lack of understanding about how *C. albicans* regulates its morphogenesis in response to these environmental changes. In addition, the synchronised morphogenesis of *C. albicans* cells, and other dimorphic fungi, seems to be coordinated within the cell population by chemical signals (Chen and Fink, 2006; Hogan, 2006a,b; Nickerson et al., 2006; Chen et al., 2007). Dimorphic fungi are known to produce several signaling metabolites. Farnesol, which suppresses filamentous formation in *C. albicans*, is the best characterized of these molecules (Hornby et al., 2001), but the mechanism by which farnesol is sensed by *C. albicans* is not yet clear. Tyrosol is another signaling molecule produced by *C. albicans* and it stimulates the yeast-to-hypha conversion (Chen et al., 2004). Other metabolites such as estradiol (mammalian metabolite) and

Table 1
The effect of environmental cues on the morphogenesis of *C. albicans*.

Environmental cues	Effect on morphogenesis	References
25 °C or lower	↑Yeast formation	Lee and Mitchell (1979)
37 °C or higher	↑Pseudohypha formation (maintains the yeast form at pH 4.5)	Lee and Mitchell (1979) and Lee et al. (1999)
5–15% CO ₂	↑Predominantly pseudohypha formation	Bahn and Mühlischlegel (2006), Klengel et al. (2005), Mock et al. (1990) and Sims (1986)
Ca ²⁺	↑Hypha formation ↓Hypha formation (absence of other divalent ions)	Brown et al. (1999)
D-glucose	↑Hypha formation (in presence of Mg ²⁺) ↑Hypha formation (at 37 °C and pH 7–8)	Holmes et al. (1991) and Sabie and Gadd (1989)
0.2% of glucose, galactose, fructose or sucrose	↑Hypha formation	Hrmova and Drobnica (1981), Hudson et al. (2004) and Vidotto et al. (1996a)
Fructose	↑Hypha formation (in the absence of nitrogen source)	Maidan et al. (2005b)
Lee's medium	↑Hypha formation (at 37 °C) ↓Hypha formation (at 25 °C)	Vidotto et al. (1996b)
Lithium	↓Hypha formation (except in liquid culture and requires at least 15 mM LiCl)	Lee et al. (1975)
Low ammonia medium	↑Hypha formation (at pH 6.7)	Martins et al. (2008)
N-acetylglucosamine	↑Hypha formation (inhibited when pH < 4.5)	Eisman et al. (2006) and Holmes and Shepherd (1987)
pH 4.5 (acid pH)	↑Yeast formation (in the presence of glucose)	Holmes and Shepherd (1987) and Sullivan and Shepherd (1982)
pH 6–8	↑Pseudohypha formation (but maintains the yeast form at 25 °C)	Lee and Mitchell (1979) and Pollack and Hashimoto (1987)
Proline	↑Hypha formation (not observed when pH < 5.0)	Dabrowa et al. (1976) and Holmes and Shepherd (1987)
Serum	↑Hypha formation	Hilmioglu et al. (2007) and Reynolds and Braude (1956)
Spider medium	↑Hypha formation	Liu et al. (1994) and Toenjes et al. (2005)
3-oxo-C12-homoserine lactone	↓Hypha formation ↓Pseudohypha formation	Hogan et al. (2004)
Dodecanol	↓Hypha formation ↓Pseudohypha formation	Hogan et al. (2004)
Human blood without white blood cell	↑Hypha formation	Fradin et al. (2005)
Phagocytosis or surrounded by neutrophils	↓Hypha formation	Barelle et al. (2006), Fradin et al. (2005) and Rubin-Bejerano et al. (2003)
Phagocytosis by macrophages	↑Hypha formation	Barelle et al. (2006), Fradin et al. (2005), Lorenz et al. (2004) and Rubin-Bejerano et al. (2003)
Parenteral lipid emulsion	↑Germ tube formation	Swindell et al. (2009)

Note: ↑: inducing, ↓: suppressing




Fig. 1. Signal transduction pathways and transcriptional regulators that affect the filamentous growth of *C. albicans* and the effect of farnesol on some of these pathways. In response to filamentous-inducing conditions, GTPases (Gpa2p, Ras1p, and Cdc42p) activate two well-characterized signaling pathways; the Cdk1p mediated MAP kinase pathway and cAMP-PKA pathway. These lead to the activation of transcriptional regulators Cph1p and Efg1p, respectively, which promote filamentous growth. Nitrogen starvation activates both the MAP Kinase and cAMP-PKA pathways via ammonium permease (Mep2p). Adenylyl cyclase (Cdc35p) not only responds to Ras1p, it is also activated in response to G-proteins (Gpr1p and Gpa2p), which are activated by glucose deficiency and the presence of methionine. Cdc35p also acts as a sensor for CO₂ and peptidoglycans. Oxidative stress and osmotic stress are sensed by a two component system (Sln1p and Ssk1p), which in turn suppresses Hog1p MAP kinase pathways. At 37 °C, PI4p5kp synthesizes phosphatidylinositol 4,5-bisphosphate (PIP₂) to promote the release of intracellular Ca²⁺. The role of other individual genes are summarized in Table 2. Quorum sensing molecule farnesol inhibits filamentous growth by suppressing the expression of RAS1, SHO1, HST7, CEK1, and HWP1, while it upregulates filamentous-suppressing genes such as TUP1 and HOG1. Blue boxes represent the membrane-anchored proteins, red boxes represent the transcriptional regulators, and the purple box indicates DNA-binding proteins (Nrg1p, Mig1p, or Rfg1p). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

dibutyryl cAMP are also known to affect *C. albicans* morphogenesis (Niimi et al., 1980; White and Larsen, 1997; Cheng et al., 2006). Despite many studies showing the effect of small molecules on fungal morphogenesis, the pathways that recognize such signals and their effect on virulence are poorly characterized. Thus, this article reviews our current knowledge of *C. albicans* morphogenesis from a metabolic point of view and relates it to the activity of central carbon metabolism and the biosynthesis of quorum sensing molecules.

2. Signaling pathways and transcriptional regulators associated with *C. albicans* morphogenesis

In the last 10 years, the use of gene knockout mutagenesis and transcriptional studies have revealed signaling pathways, transcriptional factors, as well as other regulatory components that collectively drive the yeast-to-hyphal transition, or its reversion (Gow, 2009). The roles of signaling pathways and transcriptional regulators on *C. albicans* morphogenesis have been discussed in

many review articles (Alonso-Monge et al., 2009b; Bahn et al., 2007; Biswas et al., 2007; Brown and Gow, 1999; Dhillon et al., 2003; Enrst, 2000; Hogan and Sundstrom, 2009; Lengeler et al., 2000; Liu, 2001, 2002; Navarro-García et al., 2001; Román et al., 2007; Wang, 2009). An important class of proteins involved early in signaling pathways is the GTPase superfamily. These G-proteins bind and hydrolyse guanosine triphosphate (GTP) in response to certain environmental stimuli (e.g. glucose, proline, pH, nitrogen deficiency, serum, and oxidative stress), and regulate downstream signal transduction. In *C. albicans*, the GTPases that regulate morphogenesis include G-protein-coupled receptor Gpr1p (Maidan et al., 2005), and the following members of the Ras superfamily: Rho-type Cdc42p (Johnson, 1999); RHeb-type Rhb1p (Tsao et al., 2009); and a small GTPase, Ras1p (Feng et al., 1999; Leberer et al., 2001).

GTPases are thought to be the main regulators that activate the two best-studied signaling pathways involved in *C. albicans* morphogenesis: the cAMP-Protein Kinase A (PKA), and the Mitogen-Activated Protein Kinase (MAPK), pathways (Fig. 1). The cAMP-PKA pathway involves adenylyl cyclase (Cdc35p) (Rocha et al.,

Table 2Genes involved in the morphogenesis of *C. albicans*.

Genes	Gene functions	Involvement in morphogenesis	References
<i>Signaling pathway genes</i>			
<i>BCY1</i>	The regulatory subunit of PKA	↑Hypha formation	Cassola et al. (2004)
<i>CAP1</i>	Adenylate cyclase-associated protein	↑Hypha formation	Bahn and Sundstrom (2001)
<i>CDC42</i>	Rho-type GTPase; a main regulator of cell polarity in fungi with two forms; CDC42-GTP and CDC42-GDP. CDC42-GTP activates <i>CST20</i> and promotes septin ring formation	Only CDC42-GTP: ↑hypha formation	Bassilana et al. (2003), Su et al. (2005) and Court and Sudbery (2007)
<i>CDC24</i>	The GDP-GTP exchange factor for <i>CDC42</i> ; promotes CDC42-GTP formation	↑Hypha formation	Bassilana et al. (2003)
<i>RGA2, BEM3</i>	CDC42 GTPase-activating proteins; promotes CDC42-GDP formation	↓Hypha formation	Court and Sudbery (2007) and Wang (2009)
<i>CDC35 or CYR1</i>	Adenylyl cyclase, a sensor that responds to CO ₂ and bacterial peptidoglycan (PGN)-like molecules	↑Hypha formation	Rocha et al. (2001) and Klengel et al. (2005)
<i>CDC53</i>	Encodes cullin, part of ubiquitin-ligase complex SCF	↑Pseudohypha formation	Xu et al. (2008)
<i>CST20</i>	MAPKK kinase of <i>CEK1</i> pathway	↓hypha formation	Trunk et al. (2009)
<i>HST7^a</i>	MAPK kinase of <i>CEK1</i> pathway	↑Hypha formation	Leberer et al. (1996)
<i>CEK1^a</i>	MAP kinase of <i>CEK1</i> pathway; phosphorylation is prevented by farnesol	↑Hypha formation	Leberer et al. (1996)
<i>CPP1</i>	Cek1p phosphatase, inactivates the Cek1p	↑Hypha formation	Csank et al. (1998) and Román et al. (2009)
<i>CRK1^a</i>	Cdc2-related kinase, suppressed by farnesol	↓Hypha formation	Csank et al. (1997)
<i>GPA2</i>	A G-protein α subunit	↑Hypha formation	Chen et al. (2000) and Sato et al. (2004)
<i>MKC1</i>	MAPK kinase; maintains cell wall integrity	↑Hypha formation	Sánchez-Martínez and Pérez-Martín (2002) and Maidan et al. (2005)
<i>YPD1</i>	A phosphohistidine intermediate protein that transfers a phosphate from Sln1p to Ssk1p	Expressed in both yeast and hyphal forms	Navarro-García et al. (1998)
<i>SSK1</i>	Response regulator of two component system; suppresses <i>HOG1</i> -mediated MAPK pathways and it is associated with cell wall synthesis and oxidative-stress response	Payne et al. (2000)	Calera et al. (2000) and Menon et al. (2006)
<i>SSK2</i>	MAPKK Kinase of <i>HOG1</i> pathway	↑Hypha formation	Cheetham et al. (2007)
<i>PBS2</i>	MAPK Kinase of <i>HOG1</i> pathway, essential for oxidative-stress response	↓Hypha formation	Arana et al. (2005)
<i>HOG1^a</i>	High osmolarity glycerol MAP kinase; responds to osmotic stress, temperature upshift, oxidative stress. Represses the activity of <i>CEK1</i> , regulates respiratory metabolism, and upregulated by farnesol	↓Hypha formation	José et al. (1996) and Eisman et al. (2006)
<i>PDE2^a</i>	Phosphodiesterase, suppressed by farnesol	Alonso-Monge et al. (2003), Smith et al. (2004) and Alonso-Monge et al. (2009a)	↑Hypha formation
<i>PI4P5 K</i>	phosphatidylinositol-4 phosphate 5-kinase; synthesizes phosphatidylinositol 4,5-bisphosphate (PIP ₂) in response to high temperature	↑Pseudohypha formation	Bahn et al. (2003) and Sato et al. (2004)
<i>RAD53</i>	The kinase involved in mediating DNA damaged-induced hyphal growth	↑Hypha formation	Smith et al. (2004)
<i>RAS1^a</i>	A member of the GTPase superfamily; responds to nitrogen starvation, glucose and serum. Suppressed by farnesol	↑Hypha formation	Hairfield et al. (2002)
<i>CDC25</i>	<i>RAS1</i> guanine exchange factor; responds to glucose and induces cAMP signaling pathway	↑Hypha formation	Shi et al. (2007)
<i>RHB1</i>	Rheb of Ras superfamily, involves in the nitrogen starvation inducing filamentous growth	↑Hypha formation	Feng et al. (1999)
<i>TSC2</i>	Homolog of human tuberous sclerosis protein 2, negatively regulates the GTPase activity of <i>RHB1</i>	↑Hypha formation	Enloe et al. (2000)
<i>SHO1</i>	An adaptor protein that responds to oxidative stress and cell wall biosynthesis. It activates <i>CEK1</i> Kinase pathway.	↓Hypha formation	Tsao et al. (2009)
<i>TPK1</i>	The catalytic subunit of PKA	↑hypha formation	Tsao et al. (2009)
<i>TPK2</i>	The catalytic subunit of PKA	↑Hypha formation	Román et al. (2005)
<i>YAK1</i>	Ser/Thr protein kinase; Upstream of Tup1p	↑Hypha formation	Cloutier et al. (2003)
<i>Transcription factor genes</i>			
<i>CPH1</i>	Transcription factor that is activated by MAPK pathway; responds to starvation and GlcNAc	↑Hypha formation	Cloutier et al. (2003)
<i>CZF1</i>	Putative zinc finger transcription factor; responds to agar-embedded growth conditions	↑Pseudohypha formation	Goyard et al. (2008)
<i>EFG1</i>	Transcription factor that is activated by PKA pathway; responds to serum, upregulates glycolysis and downregulates TCA cycle	↑Hypha formation	Lo et al. (1997)
<i>TEC1</i>	A transcription factor - member of the TEA/ATTS family; regulated by <i>EFG1</i>	↑Pseudohypha formation	Brown et al. (1999)
<i>TUP1^a</i>	A general transcriptional repressor; regulates glycolysis and TCA cycle. Upregulated by farnesol	↑Hypha formation	Stoldt et al. (1997) and Leng et al. (2001)
<i>SSN6</i>	A general transcriptional repressor forms a complex with <i>TUP1</i>	↓Hypha formation	Schweizer et al. (2000) and Lane et al. (2001)
<i>MIG1</i>	DNA-binding protein (DBP) binds to <i>TUP1</i>	↓Pseudohypha formation	Braun and Johnson (1997), Cao et al. (2005) and Kebaara et al. (2008)
<i>NRG1</i>	DBP binds to <i>TUP1</i>	↓Hypha formation	Hwang et al. (2003)
		↓Pseudohypha formation	Murad et al. (2001)
			Braun et al. (2001) and Murad et al. (2001)

Table 2 (continued)

Genes	Gene functions	Involvement in morphogenesis	References
<i>RFG1</i>	Repressor for filamentous growth; DBP binds to <i>TUP1</i>	↓Hypha formation	Kadosh and Johnson (2001) and Khalaf and Zitomer (2001)
<i>RIM101</i>	pH-response regulator, activated by alkaline conditions and regulates both <i>PHR1</i> and <i>PHR2</i> . It is required for ARO9 pH dependent expression and required for virulence during systemic infection	↓Pseudohypha formation ↑Hypha formation	Ramon et al. (1999) and Nobile et al. (2008)
<i>RGT1</i>	A transcriptional repressor for genes that encode hexose transporters, N-acetylglucosamine transporter (<i>NGT1</i>), and downregulates glycolysis	↓Hypha formation	Sexton et al. (2007)
<i>GCN4</i>	General amino acid control; regulates both morphogenesis and metabolic responses to amino acid starvation	↑Hypha formation	Tripathi et al. (2002)
<i>UME6</i>	A transcriptional regulator; encodes zinc finger DNA-binding protein	↑Pseudohypha formation ↑Hypha formation	Strich et al. (1994) and Banerjee et al. (2008)
<i>Receptor genes</i>			
<i>CDR1, CDR2</i>	Multidrug transporters of the ABC family; an efflux pump that removes estradiol from the cell	↑Germ tube is formed in the present of estradiol	Cheng et al. (2006)
<i>GAP1</i>	General amino acid permease; the expression of transporter is induced by GlcNAc and Cph1p.	↑Hypha formation	Biswas et al. (2003)
<i>HGT4</i>	Glucose and galactose sensor	↑Hypha formation	Brown et al. (2006) and Brown et al. (2009)
<i>GPR1</i>	G-protein-coupled receptor; upstream of cAMP-PKA pathway, senses methionine and proline	↑Hypha formation	Maidan et al. (2005) and Maidan et al. (2005)
<i>HGT4</i>	Glucose sensor; induces the expression of hexose transporters (e.g. <i>HGT12</i> , <i>HXT12</i> and <i>HGT7</i>)	↓Hypha formation	Brown et al. (2006)
<i>CHK1^a/NIK1/ SLN1</i>	Histidine kinase of two component system; <i>SLN1</i> is the upstream components of Hog1-mediated MAPK pathway. The <i>CHK1</i> mutant overcomes the hyphal-suppressing effect of farnesol	↓Pseudohypha formation ↑Hypha formation	Nagahashi et al. (1998) and Calera and Calderone (1999), Yamada-Okabe et al. (1999) and Kruppa et al. (2004)
<i>NGT1</i>	N-acetylglucosamine transporter	In a homozygous <i>NGT1</i> deletant, hyphal growth cannot be induced by GlcNAc	Alvarez and Konopka (2007)
<i>MEP2</i>	Ammonium permease; senses nitrogen starvation. Expression is suppressed by: 10 mM of NH_4^+ , overexpression of <i>RHB1</i> , and <i>TSC2</i> deletion. It activates both MAP kinase pathway and cAMP-PKA pathway	↑Pseudohypha formation	Biswas and Morschhäuser (2005) and Tsao et al. (2009)
<i>Metabolic genes</i>			
<i>ALO1</i>	D-arabinono-1,4-lactone oxidase; catalyzes the last step of D-erythroascorbic acid biosynthesis	↑Hypha formation	Huh et al. (2001)
<i>ATC1</i>	Cell wall-linked acid trehalase, converts exogenous galactose into glucose.	↑Hypha formation	Pedreño et al. (2007)
<i>GAL10</i>	UDP-galactose-4-epimerase; converts galactose into glucose	↑Pseudohypha formation ↓Hypha formation	Singh et al. (2007)
<i>PDX1</i>	Pyruvate dehydrogenase	↑Hypha formation ↑Pseudohypha formation	Vellucci et al. (2007)
<i>TSP1</i>	Trehalose-6-phosphate synthase; involved in trehalose biosynthesis.	↑Hypha formation	Zaragoza et al. (1998)
<i>Cell wall genes</i>			
<i>ALS1</i>	Adhesin; hypha specific gene	↑Hypha formation ↑Pseudohypha formation	Fu et al. (2002)
<i>HWP1^a</i>	Cell wall protein; hyphal-specific gene, involved in cellular adhesion during biofilm formation, and suppressed by farnesol	↑Hypha formation	Sharkey et al. (1999) and Ramage et al. (2002), Enjalbert and Whiteway (2005) and Padovan et al. (2009)
<i>INT1</i>	Integrin-like, role in adhesion	↑Hypha formation ↑Pseudohypha formation	Gale et al. (1998) and Asleson et al. (2001)
<i>PHR1</i>	GPI-anchored glycosidase; cell wall structure, expressed at pH 5.5 or alkaline pH	↑Hypha formation	Ghannoum et al. (1995) and De Bernardis et al. (1998)
<i>PHR2</i>	GPI-anchored glycosidase; cell wall structure, expressed at acidic pH	↑Pseudohypha formation ↑Hypha formation	Mühlschlegel and Fonzi (1997) and De Bernardis et al. (1998)
<i>PMT1</i>	Mannosyltransferase	↑Hypha formation	Timpel et al. (1998)
<i>PMT6</i>	Mannosyltransferase, role in adhesion and antifungal resistance	↑Hypha formation	Timpel et al. (2000)
<i>Cell cycle-associated genes</i>			
<i>CDC28</i>	Cyclin-dependent kinase; phosphorylates <i>Cdc11</i> p(septin gene) in association with either <i>CCN1</i> or <i>HGC1</i> . <i>CDC28-HGC1</i> negatively regulates <i>RGA2</i> and also phosphorylates Efg1p	↑Hypha formation	Sinha et al. (2007) and Wang et al. (2009)
<i>HGC1</i>	Hyphal-specific G1 cyclin related protein; maintains the localization of polarity proteins (e.g. actin) on hyphal tip, regulated by cAMP-PKA pathway and Tup1p	↑Hypha formation	Zheng and Wang (2004)
<i>CCN1</i>	G ₁ cyclin; role in maintaining hyphal growth	↑Hypha formation	Loeb et al. (1999)

(continued on next page)

Table 2 (continued)

Genes	Gene functions	Involvement in morphogenesis	References
<i>CLN3</i>	G ₁ cyclin; is important for yeast budding and negatively regulates the yeast-to-hyphal transition	↓Hypha formation	Bachewich and Whiteway (2005) and Lazo et al. (2005)
<i>CLB2, CLB4</i>	B-cyclins that are negative regulators of polarized growth	↓Hypha formation ↓Pseudohypha formation	Bensen et al. (2005)
<i>Others genes</i>			
<i>CDC10, CDC11</i>	Non-essential septin genes	<i>cdc10</i> and <i>cdc11</i> mutants show abnormal hyphal structure	Warenda et al. (2003)

Note: ↑: induction, ↓: suppression.

Observations are based on knockout mutagenesis studies.

^a The expression of these genes is regulated by farnesol.

2001) and PKA (Bcy1p, Tpk1p and Tpk2p) (Cassola et al., 2004; Cloutier et al., 2003) (Table 2, Fig. 1). Eventually, the PKA pathway activates an important transcription factor encoded by *EFG1* (Stoldt et al., 1997) that induces the expression of hyphal-specific genes (e.g. *HWPI*, *HYR1* and *ALS1*) (Fu et al., 2002; Leng et al., 2001; Sharkey et al., 1999). MAPK pathways typically consist of three kinases; MAPKKK, MAPKK, and MAPK that are sequentially activated one after the other by phosphorylation. There are several different MAPK pathways that have been described and linked to the morphological changes of *C. albicans* (e.g. *Cek1p*, *Hog1p*, and *Mkc1p* MAPK pathways) (Alonso-Monge et al., 2009b; Cheetham et al., 2007; Csank et al., 1998; Eisman et al., 2006; Leberer et al., 1996, 2001; Navarro-García et al., 1998). Fig. 1 illustrates how these signaling pathways and regulatory components are interconnected, and Table 2 summarizes how these components lead to modifications in cell wall organization, cell polarity, metabolism, cell cycle, virulence factors, and morphogenesis. Furthermore, transcriptome studies have identified several hyphal-suppressing genes in *C. albicans*. These repressor genes, such as *TUP1*, *SSN6*, *NRG1*, *MIG1*, and *HOG1*, are involved in the regulation of central carbon metabolism (Braun and Johnson, 1997; Hwang et al., 2003; José et al., 1996; Murad et al., 2001) as illustrated in Table 3. There are many other genes that are known to be involved in the morphological switch of

C. albicans. The role of those genes and how they may influence, or have their expression changed during, morphogenesis are briefly summarized in Table 2.

3. Quorum sensing molecules and *C. albicans* morphogenesis

The phenomenon of “quorum sensing” is being increasingly recognized as a fundamental aspect of microbial cell-to-cell communication and signaling. Fuqua et al. (1994) first coined this term to describe the cooperative behaviour of bacterial cells that can only take place when a certain cell population density threshold is reached. Quorum sensing is often referred to as autoinduction, a process by which individual cells release small diffusible molecules into their environment and these molecules are sensed by all cells in the population (Gray et al., 1994; Hense et al., 2007; Nealson, 1977). When high cell density is reached, these autoinducing molecules accumulate above a certain threshold level, activating and/or repressing certain genes (Fuqua et al., 1994), which in turn induce complex cellular behaviour such as secretion of extracellular enzymes (Rosenberg et al., 1977), bioluminescence (Eberhard, 1972; Eberhard et al., 1981; Fuqua et al., 1994), plasmid transfer (Piper et al., 1993), antibiotic biosynthesis (Bainton et al.,

Table 3

The effect of morphogenesis-related genes, and farnesol, on *Candida albicans* metabolism based on DNA array studies or large-scale protein analyses^a.

Metabolism of:	<i>EFG1</i>	<i>GCN4</i>	<i>HOG1</i>	<i>TUP1</i>	<i>MIG1</i>	<i>NRG1</i>	<i>RIM101</i>	<i>SSN6</i>	<i>NGT1</i>	Farnesol
Alanine										
Arginine		↑					↑pH8			
Asparagine		↑					↓pH4			
Aspartate										
Cysteine										
Ergosterol biosynthesis	↓					↓				
Glutamate	↓	↑		↓	↓	↓	↑pH8			
Glutamine										
Glycine										↑
Glycolysis	↑	↑	↑					↓	↓	↓
Glyoxylate shunt										↓
Histidine		↑		↓			↓pH4 ↑pH8 ↓pH8			
Isoleucine		↑								
Leucine		↑		↓	↓	↓				
Lysine		↑								
Methionine		↑								
Pentose phosphate		↑								
Phenylalanine		↑								
Proline		↑								
Serine		↑								
TCA cycle	↓	↑	↓							
Threonine		↑								
Tryptophan		↑								
Tyrosine		↑								
Valine		↑								

Note: ↑: gene/farnesol upregulates one or more enzymes in a particular biochemical pathway.

↓: gene/farnesol downregulates one or more enzymes in a particular biochemical pathway ↓^{pH4}: assayed at pH 4, ↑^{pH8}: assayed at pH 8.

1992), biofilm development (Alem et al., 2006), and morphological switches (Hornby et al., 2001). Despite quorum sensing being well-characterized and evolutionarily-conserved between diverse bacterial species (Gray and Garey, 2001; Gray et al., 1994), only a few quorum sensing like-responses have been reported in eukaryotes, such as *H. capsulatum* (Kügler et al., 2000), *C. albicans* (Chen et al., 2004; Hornby et al., 2001), and *Saccharomyces cerevisiae* (Chen and Fink, 2006). Curiously, in each of these studies quorum sensing was associated with the morphological switch from yeast-to-filamentous forms, or vice versa.

Eukaryotic quorum sensing is best studied in *C. albicans*. This fungus has the greatest number of quorum sensing molecules (QSMs) identified to date. Tryptophol and phenylethyl alcohol were the first QSMs identified in *C. albicans* (Lingappa et al., 1969). These QSMs are produced by *C. albicans* and inhibit both cell growth and germ tube formation – whether the inhibition of germ tube formation is due to the inhibition of growth remains unclear. In recent years, three additional QSMs have been isolated from *C. albicans*; farnesol, tyrosol and farnesoic acid (Chen et al., 2004; Hornby et al., 2001; Hornby and Nickerson, 2004; Oh et al., 2001), although farnesoic acid has only been reported in one *C. albicans* strain (ATCC 10231, Oh et al., 2001) and has a lower activity than farnesol (Hornby and Nickerson, 2004).

Farnesol is the best characterized QSM in *C. albicans* and it is known to block the yeast-to-filamentous transition at high cell density, as well as under other hyphal-inducing conditions (e.g. serum, proline, and N-acetylglucosamine) (Hornby et al., 2001); but it is incapable of stopping the elongation of pre-existing hyphae (Mosel et al., 2005; Navarathna et al., 2005). Moreover, farnesol is continuously released into the environment during growth, and its accumulation in the medium is roughly proportional to the cell density (Hornby et al., 2001). It has been proposed that farnesol is secreted by the cells to inhibit germ tube formation in the late stage of biofilm development where there is a high density of interwoven filamentous cells, and, therefore, it promotes the dispersal of yeast cells to colonize new environments (Alem et al., 2006). After farnesol was discovered in *C. albicans* cultures, several studies have been undertaken to investigate the mechanisms by which QSMs affect morphogenesis. For instance, Sato et al. (2004) used RT-PCR to show that farnesol inhibited MAP kinase cascades via the suppression of *HST7* and *CPH1* gene expression (Fig. 1). Hog1p phosphorylation also increased in the presence of farnesol (Smith et al., 2004). Cao et al. (2005) demonstrated that several morphogenesis-associated genes were downregulated (e.g. *CRK1* and *PDE2*) and some upregulated (e.g. *TUP1*) by the presence of farnesol. It has been suggested that hyphal formation induced by the cAMP-PKA pathway can be repressed by farnesol which suppresses the *RAS1*-*CDC35* pathway (Davis-Hanna et al., 2008). Kebara et al. (2008) showed that farnesol inhibits hyphal formation by upregulating the global repressor of *TUP1* because it failed to suppress hyphal development in *tup1/tup1* and *nrg1/nrg1* null-mutants. Recently, Román et al. (2009) showed that farnesol prevents the activation of Cek1p, which is a part of the MAPK cascade. In summary, the studies to date show that farnesol inhibits *C. albicans* morphogenesis by suppressing both MAPK and cAMP-PKA pathways and promoting the expression of hyphal-suppressor genes such as *TUP1* and *HOG1* (Fig. 1). How yeast cells sense farnesol, however, is still unclear.

Tyrosol is another QSM produced by *C. albicans*. It has been observed that when a dense culture of *C. albicans* is diluted into fresh medium there is long lag-phase before exponential growth initiates (Chen et al., 2004). This lag period can be shortened, and germ tube formation can be enhanced, when tyrosol is supplied (Chen et al., 2004). Alem et al. (2006) also suggested that tyrosol stimulated hypha formation in the early and intermediate phases of *C.*

albicans biofilm formation. It seems that tyrosol does not induce germ tube formation when *C. albicans* grows under non-filamentous-inducing conditions but rather accelerates the morphological switch from yeast-to-hyphal growth under favourable environmental conditions (Chen et al., 2004). In contrast to farnesol, little is known about how tyrosol exerts its effects. A study of gene expression profiles of *C. albicans* cells at different cell densities with and without tyrosol suggested that the acceleration of germ tube formation may simply be due to upregulation of genes associated with DNA replication (e.g. DNA polymerase, chromosome-separation factor) and the cell cycle (e.g. cell cycle checkpoint protein) (Chen et al., 2004).

Together, these studies demonstrate that the morphogenesis of *C. albicans* is under the positive and negative regulation of QSMs in response to cell density. QSMs such as tryptophol, phenylethyl alcohol, and farnesol suppress hyphal formation when cells grow at high density. The QSM tyrosol accelerates germ tube formation at low cell density (Fig. 2). QSMs appear to have other roles in addition to affecting morphogenesis. Farnesol, for example, enhances *C. albicans* resistance to oxidative stress (Westwater et al., 2005), and tyrosol can act as an antioxidant, protecting *C. albicans* cells during phagocytosis by neutrophils (Cremer et al., 1999). Moreover, when growing in the human body *C. albicans* often grows in a polymicrobial environment and QSMs appear to be also involved in inter-species competition (Shank and Kolter, 2009). Farnesol not only inhibits filament formation in other *Candida* species (e.g. *Candida dubliniensis* and *Candida tropicalis*) (Henriques et al., 2007; Zibafar et al., 2009), it also induces apoptosis in some fungi (e.g. *S. cerevisiae*, *Aspergillus nidulans*, and *Penicillium expansum*), and suppresses the growth of *Paracoccidioides brasiliensis* (Derengowski et al., 2009; Fairn et al., 2007; Liu et al., 2009; Semighini et al., 2006). In contrast, when *C. albicans* is exposed to other QSMs such as 3-OXO-C12 homoserine lactone from *Pseudomonas aeruginosa*, its filamentous growth is suppressed (Hogan et al., 2004). Interestingly, farnesol seems to be also involved in *C. albicans*-host interactions. Proteomic approaches have shown that farnesol triggers apoptosis in both human oral carcinoma cells and *C. albicans* itself via classic apoptotic pathways (Scheper et al., 2008; Shirtliff et al., 2009). It appears that *C. albicans* is more susceptible to farnesol-mediated cell death when log-phase cells grow under nutrient-poor conditions (Langford et al., 2010). Dècanis et al. (2009) demonstrated that farnesol upregulates the expression of toll-like receptor 2, and increases the production of interleukin-6 and β -defensin 2 in the engineered tissue of human oral mucosa. The same authors suggest that farnesol promotes epithelial cell immunity against *C. albicans* (Dècanis et al., 2009). Moreover, Ghosh et al. (2010) showed that farnesol induces the expression of inflammatory cytokines in the macrophage. Therefore, these studies clearly demonstrate that QSMs play complex roles in *C. albicans*-host and *C. albicans*-interspecies interactions.




Fig. 2. The effect of quorum sensing molecules on the dimorphic transition of *C. albicans*. Tyrosol accelerates hypha formation, while farnesol, tryptophol, and phenylethyl alcohol suppress hypha development.

4. *In silico* reconstruction of the central carbon metabolism of *C. albicans* and the biosynthesis of quorum sensing molecules

The central carbon metabolism of most organisms is highly conserved comprising both catabolic and anabolic biochemical reactions. Central carbon metabolism is indispensable for cellular growth and any major cellular event, such as morphogenesis, will certainly be accompanied by significant changes in these important pathways. Therefore, knowledge of the different biochemical pathways involved in the central carbon metabolism of *C. albicans*, as well as how they are affected by the morphogenetic process, is crucial to understand the mechanisms of morphogenesis in this fungus.

Most of the genes thought to encode enzymes involved in the central carbon metabolism of *C. albicans* have been annotated by individual *in vitro* studies and by high-throughput transcriptomics (Bensen et al., 2004; Doedt et al., 2004; Enjalbert et al., 2006; Fradin et al., 2005; García-Sánchez et al., 2004; García-Sánchez et al., 2005; Marcus et al., 2004; Hromatka et al., 2005; Karababa et al., 2004; Lan et al., 2004; Lorenz et al., 2004; Murad et al., 2001; Nantel et al., 2002; Nett et al., 2009; Rogers and Barker, 2002; Sexton et al., 2007; Shirliff et al., 2009; Singh et al., 2005; Swoboda et al., 1994; Tournu et al., 2005) and proteomics studies (Cabezón et al., 2009; Fernández-Arenas et al., 2007; Hernández et al., 2004; Kusch et al., 2008; Pitarch et al., 2004; Shirliff et al., 2009; Thomas et al., 2006; Urban et al., 2003; Yin et al., 2004). We have reconstructed *in silico* the central carbon metabolism of *C. albicans* directly from its non-annotated genome sequence available from GenBank, using a program called IdentiCS (Sun and Zeng, 2004). Then the metabolic pathways were curated based on three publicly available online-databases: the *Candida* Genome Database – CGD (<http://www.candidagenome.org/>), the *Saccharomyces* Genome Database – SGD (<http://www.yeastgenome.org/>), and the Kyoto Encyclopedia of Genes and Genomes-KEGG (<http://www.genome.jp/kegg/>) (Fig. 3).

4.1. Glycolysis

Genes encoding all glycolytic enzymes in *C. albicans* have been identified and are conserved across fungi. Although the mRNA expression of glycolytic genes is not regulated tightly and fluctuates during the yeast-to-hyphal switch (Swoboda et al., 1994), some differences in the activities of glycolytic enzymes have been correlated with yeast and hyphal growth (Schwartz and Larsh, 1982). For instance, the specific activity of hexokinases in the hyphal form is almost twice that in the yeast form (Schwartz and Larsh, 1982). This might indicate a higher glycolytic flux during hyphal growth. Several glycolytic genes are regulated by signaling pathways associated with morphogenesis. These include enzymes such as hexokinase II (upregulated by Efg1p and downregulated by Ssn6p and Rgt1p) (Doedt et al., 2004; García-Sánchez et al., 2005; Sexton et al., 2007), phosphofructokinase (downregulated by Rgt1p) (Sexton et al., 2007), glucose-6-phosphate isomerase (upregulated by Efg1p) (Doedt et al., 2004), fructose-bisphosphate aldolase (upregulated by Efg1p, Gcn4p, Hog1p) (Doedt et al., 2004; Enjalbert et al., 2006; Yin et al., 2004), phosphoglycerate kinase (upregulated by Gcn4p and Hog1p) (Enjalbert et al., 2006; Yin et al., 2004), and pyruvate kinase (upregulated by Gcn4p and Hog1p) (Enjalbert et al., 2006; Yin et al., 2004). In addition, Shirliff et al. (2009) used a proteomic analysis to demonstrate that several glycolytic enzymes (e.g. glyceraldehyde 3-phosphate dehydrogenase, enolase, phosphoglycerate mutase and pyruvate kinase) were downregulated when *C. albicans* was exposed to farnesol.

4.2. Pentose phosphate pathway

In anaerobic respiration, the pentose phosphate pathway is coupled with glycolysis to generate both cytosolic NADPH and ribose.

The NADPH provides oxidizing energy for biosynthetic reactions (e.g. synthesis of amino acids, fatty acids, and sugar alcohols) while ribose is used in the biosynthesis of nucleotides (e.g. nucleic acids and redox cofactors). The conversion of glucose 6-phosphate to 6-phosphogluconolactone by a NADP-dependent glucose 6-phosphate dehydrogenase (G6PDH) is the first reaction, and the key-regulatory step, in the pentose phosphate pathway. A putative G6PDH is encoded by the *ZWF1* gene in *C. albicans* and its expression is upregulated by Gcn2p and Gcn4p (Tournu et al., 2005).

4.3. Pyruvate dehydrogenase

In aerobic respiration, pyruvate produced by glycolysis and the pentose phosphate pathway is mainly oxidatively decarboxylated to acetyl-CoA by the pyruvate dehydrogenase multi-complex enzyme. This enzyme is highly conserved across species, and the *C. albicans* genome contains genes with high homology to the *S. cerevisiae* operon encoding all subunits of pyruvate dehydrogenase. Disruption of one of the pyruvate dehydrogenase subunit genes (*PDX1*) in *C. albicans* resulted in a deficiency in filamentous growth (Vellucci et al., 2007). Thus the availability of acetyl-CoA may be required for the morphogenesis of *C. albicans*.

4.4. TCA cycle

The tricarboxylic acid (TCA) cycle is a central hub of carbon metabolism (Fig. 3) and, under aerobic conditions, acetyl-CoA formed from pyruvate is oxidized completely to carbon dioxide, water, and chemical energy through this pathway. This process is catalyzed by a series of conserved enzymes. All genes involved in the TCA cycle have been found in the *C. albicans* genome and all the TCA cycle enzymes have been extracted from *C. albicans* cells (Rao et al., 1962). Some TCA cycle enzymes are regulated by signaling pathways. For example, α -ketoglutarate dehydrogenase, succinate dehydrogenase, fumarase, citrate synthase, and malate dehydrogenase are repressed by the Efg1p transcription factor (Doedt et al., 2004), which appears to be involved in hyphal formation (Leng et al., 2001; Stoldt et al., 1997). Citrate synthase is downregulated by Hog1p and cis-aconitase is upregulated by Gcn4p (Enjalbert et al., 2006). The TCA cycle is, however, *amphibiotic* because the cycle not only catabolically decarboxylates pyruvate but it also has anabolic functions. Intermediate compounds of the TCA cycle are used as precursors for the biosynthesis of building blocks for the cell (e.g. amino acids and lipids). Therefore, a constant inflow of carbon to supply intermediate compounds for amino acid and lipid biosynthesis is essential for cell growth. An example of a metabolic strategy to maintain the supply of TCA cycle intermediates is the glyoxylate shunt.

4.5. Glyoxylate shunt

The glyoxylate shunt assimilates two-carbon molecules such as ethanol and acetate into the TCA cycle; this process not only supplies TCA cycle substrates but also allows microbial growth in environments where only two-carbon compounds are available. The glyoxylate shunt can be considered as a shortcut version of the TCA cycle. The cycle bypasses two decarboxylation steps and directly converts isocitrate to malate using two enzymes: isocitrate lyase (encoded by *ICL1*) and malate synthase (encoded by *MLS1*) (Fig. 3). The regulation of the glyoxylate shunt is primarily via the activity of isocitrate lyase, which is downregulated by the presence of the quorum sensing molecule farnesol (Enjalbert and Whiteway, 2005) and is usually repressed by glucose in fungi. Although there has been no direct report on the role of the glyoxylate shunt in *C. albicans* morphogenesis, Lorenz and Fink (2001)




Fig. 3. The *in silico* reconstruction of the central carbon metabolism of *C. albicans*. Blue shading indicates the intermediate metabolites of glycolysis and the TCA cycle that act as key precursors for the biosynthesis of amino acids. The intermediates for biosynthesis of individual amino acids are differentiated by different colours and red 3D boxes represent the final amino acid products. Blue 3D boxes indicate quorum sensing molecules derived from aromatic amino acids and the putative genes encoding enzymes are in red text. The few reactions not verified by *in vitro* studies are indicated by red lines.

showed that the disruption of both *ICL1* and *MLS1* reduced *C. albicans* virulence using a mouse model of intravenous infection.

4.6. Amino acid biosynthesis

The biosynthesis of amino acids is achieved through a series of biochemical reactions that can be classified into six distinct pathways. Each amino acid pathway derives its carbon skeleton from a common precursor which is an intermediate compound of glycolysis or the TCA cycle. In *C. albicans*, the glycolytic pathway provides precursors for the biosynthesis of four amino acid families: the histidine, serine, pyruvate and aromatic families. Only the aspartate and glutamate family precursors are derived from the TCA cycle (Fig. 3).

4.6.1. Glutamate biosynthesis

Glutamate is synthesized directly from α -ketoglutarate by an NADPH-dependent glutamate dehydrogenase (GDH). This is an important reaction in central carbon metabolism because it serves to assimilate ammonia for the biosynthesis of several other amino acids and at the same time it maintains the redox equilibrium in the cell. The NADH-dependent glutamate dehydrogenase (Gdh2p) favors the catabolic deamination of glutamate to α -ketoglutarate. Therefore, Gdh2p is an important enzyme in the biosynthesis of other amino acids (e.g. aspartate, tyrosine, phenylalanine, etc.). Three glutamate dehydrogenase genes have been identified in *S. cerevisiae* but only *GDH2* and *GDH3* have orthologues in the *C. albicans* genome. NADH-dependent *GDH2* is transcriptionally repressed by Tup1p, Nrg1p and Mig1p (Murad et al., 2001). NADP-dependent *GDH3* is downregulated by Efg1p (Doedt et al., 2004), and upregulated 2-fold by Rim101p at pH 8 (Bensen et al., 2004).

4.6.2. Aromatic amino acid and quorum sensing molecule biosynthesis

The discovery that extracellular aromatic alcohols act as quorum sensing molecules during fungal morphogenesis makes aromatic amino acid biosynthesis an important component of the central carbon metabolism regarding morphogenesis. But how the aromatic alcohols are synthesized from aromatic amino acids in *C. albicans* is still unclear. In contrast, the production of aromatic alcohols in *S. cerevisiae* has been well-characterized. Aromatic alcohols are derived from aromatic amino acid catabolism via the Ehrlich pathway (Felix, 1907). This process in *S. cerevisiae* involves three consecutive enzymatic steps; a transaminase (encoded by *ARO8*, *ARO9*), a decarboxylase (encoded by *ARO10*), and reduction by alcohol dehydrogenase (encoded by *ADH*) (Hazelwood et al., 2008; Sentheshanmuganathan, 1960). It is proposed that the biosynthesis of aromatic alcohols such as tyrosol, tryptophol, and phenylethyl alcohol in *C. albicans* (Fig. 2), follows the same biosynthetic pathway as in *S. cerevisiae* (Ghosh et al., 2008). This study demonstrated that the biosynthesis of aromatic alcohols by *C. albicans* decreased when the *ARO80* gene, which encodes a transcriptional activator known to increase the expression of *Aro9p* and *Aro10p* in *S. cerevisiae*, was deleted (Ghosh et al., 2008).

4.6.3. Regulation of global amino acid biosynthesis

There are several morphology-associated genes that also regulate multiple amino acid biosynthetic pathways in response to starvation and pH changes. When amino acids are limiting growth, *C. albicans* switches from yeast to filamentous growth (Tripathi et al., 2002). This phenomenon is regulated by the transcriptional activator Gcn4p in an *EFG1*-dependent fashion (Tripathi et al., 2002). The double deletion *gcn4/gcn4* blocks amino acid starvation from inducing morphogenesis, but the *efg1/efg1* knockout mutants fail to express this *GCN4* phenotype (Tripathi et al., 2002). Gcn4p is not only involved in the dimorphic switch of *C. albicans*, but it also

acts as a global regulator of metabolism in response to amino acid starvation (Tournu et al., 2005; Tripathi et al., 2002). Tripathi et al. (2002) found from DNA microarray analysis that, under amino acid starvation conditions, Gcn4p upregulates numerous enzymes involved in amino acid biosynthesis, except those responsible for the biosynthesis of cysteine, glycine, alanine, aspartate and glutamine (Table 3) (Tournu et al., 2005). Rim101p is a pH-response regulator that is activated at alkaline pH (Li and Mitchell, 1997). In turn, Rim101p regulates the expression of pH-response regulators (Phr1p and Phr2p) involved in the pH-dependent morphogenesis of *C. albicans* (Ramon et al., 1999). Bensen et al. (2004) compared transcriptional profiles between *rim101/rim101* null-mutants and the wild type strain. They found that Rim101p upregulates arginine, glutamine, and histidine biosynthetic enzymes at pH 8, while asparagine and histidine biosynthetic enzymes are downregulated at pH4, respectively (Table 3).

The *in silico* reconstruction of central carbon metabolic pathways based on the *C. albicans* genome (Fig. 3) enables us to integrate post-genomic studies into a metabolic network that helps elucidate the physiological and biochemical mechanisms governing the morphological switch of *C. albicans*.

5. Central carbon metabolism, virulence, and morphogenesis of *C. albicans*

In addition to its role in morphogenesis, central carbon metabolism has increasingly been recognized for its importance in fungal pathogenicity. It has been shown that the deletion of glycolytic transcriptional regulators (*TYE7*, *GAL4*), a gluconeogenic gene (*FBP1*), glyoxylate cycle genes (*ICL1*, *MLS1*), or a β -oxidation gene (*FOX2*), attenuates the virulence of *C. albicans* in a murine model of systemic infection (Askew et al., 2009; Lorenz and Fink, 2001; Piekarska et al., 2006; Ramírez and Lorenz, 2007). Interestingly, *C. albicans* appears to modulate these carbon assimilatory pathways during infection. When glucose is at similar levels to that found in the bloodstream there is an activation of glycolysis and repression of both the glyoxylate cycle and gluconeogenesis in *C. albicans* (Barelle et al., 2006; Rodaki et al., 2009). In contrast, several transcriptomic studies have demonstrated that following the phagocytosis of *C. albicans* by neutrophils or macrophages, there is a downregulation of glycolysis, and an upregulation of the glyoxylate cycle, gluconeogenesis, and β -oxidation (Fradin et al., 2005; Lorenz et al., 2004; Rubin-Bejerano et al., 2003). Since all these metabolic pathways involve the utilization of non-fermentable carbon sources (e.g. amino acids, ethanol, acetate, and fatty acids), it seems likely that *C. albicans* can alter its central carbon metabolism to utilize alternative carbon sources in response to host defences or to changes in the environment, and this metabolic reprogramming is important for the virulence of *C. albicans*. Although the morphological switch from yeast-to-hyphae is observed after *C. albicans* is engulfed by macrophages, Lorenz et al. (2004) suggested that those underlying metabolic changes are independent from morphogenesis.

Lorenz et al. (2004) also demonstrated that genes for L-arginine biosynthesis and degradation were upregulated after *C. albicans* cells were engulfed by macrophages, and other groups have shown that concentrations of CO₂ greater than 50% (v/v) enhanced filamentous growth (Bahn and Mühlischlegel, 2006; Klengel et al., 2005; Mock et al., 1990; Sims, 1986). Ghosh et al. (2009) have proposed a mechanism, involving arginine, which connects these two observations. They suggest that arginase (Car1p) converts arginine to urea, which in turn is cleaved into NH₃ and CO₂ by the amidolyase enzyme (Dur1,2p [sic]). They show that the *dur1,2/dur1,2* null-mutant cannot form germ tubes in macrophages, and that the phenomenon of arginine-induced hyphal formation is not

observed in *efg1/efg1* strains. Therefore, once inside a macrophage, it is proposed that *C. albicans* produces arginine and arginine degradation produces CO₂. The CO₂ may then activate adenylyl cyclase (Cdc35p), which in turn activates the cAMP-PKA signaling pathway and Efg1p to induce the yeast-to-hyphal transition (Fig. 1) and thereby *C. albicans* escapes from the macrophage (Klengel et al., 2005).

6. The effect of metabolic genes on the utilization of carbon sources

There are several metabolic genes associated with morphogenesis or virulence that seem to affect the ability of *C. albicans* to utilize carbon sources. Firstly, trehalose-6-phosphate synthase (*TPS1*) is a gene involved in the first step of trehalose biosynthesis and it regulates glucose influx (Ernandes et al., 1998). The *tps1/tps1* null-mutant is unable to grow on glucose or fructose as a sole carbon source but this mutant is capable of growing on other carbon sources such as galactose or glycerol (Zaragoza et al., 1998). In addition, this mutant is impaired in hyphal formation (Zaragoza et al., 1998). Another gene related to trehalose biosynthesis is the cell wall-linked acid trehalase (*ATC1*) (Pedreño et al., 2004). The *atc1/atc1* null-mutant is incapable of growing on exogenous trehalose as sole carbon source and the mutant has diminished capacity to form hyphae in various media (e.g. serum, spider medium and Lee's medium) (Pedreño et al., 2007).

Galactose metabolism also plays a role in the morphogenesis of *C. albicans*. UDP-galactose-4-epimerase (*GAL10*) is a key enzyme in galactose metabolism that converts UDP-galactose into UDP-glucose. The glucose generated can then be fed into glycolysis as a carbon and energy source. The *gal10/gal10* null-mutant is unable to grow on galactose as sole carbon source (Singh et al., 2007). Compared to wild type, the mutant also exhibits increased hyphal formation in rich media, Lee's medium and spider medium (Singh et al., 2007). Recently, Hgt4p has been reported as a galactose and glucose sensor. Indeed, the *hgt4/hgt4* null-mutant cannot grow on either glucose or galactose in the presence of the respiration inhibitor antimycin A (Brown et al., 2009, 2006). The *hgt4/hgt4* mutant is hypo-filamentous and less virulent in a mouse model of disseminated candidiasis (Brown et al., 2006).

In addition, carnitine biosynthesis and acetyl-CoA metabolism can affect the ability of *C. albicans* to utilize different carbon sources. Carnitine is an essential metabolite that acts as a shuttle to transport fatty acids and acetyl groups between intracellular compartments. Mutants lacking carnitine synthetic enzymes (e.g. trimethyl-lysine dioxygenase, trimethylaminobutyraldehyde dehydrogenase, and butyrobetaine dioxygenase) or lacking carnitine acetyltransferase, which transfers the acetyl group from acetyl-CoA to carnitine, are unable to utilize non-fermentable carbon sources such as fatty acids, acetate, or ethanol as sole carbon sources (Strijbis et al., 2009; Zhou and Lorenz, 2008). Moreover, Carman et al. (2008) have shown that a *C. albicans* mutant strain with acetyl-CoA synthase (*ACS2*) deleted was unable to utilize glucose, acetate, or ethanol but the mutant cells were viable when given fatty acids or glycerol. The same authors deleted acetyl-CoA hydrolase (*ACH1*), an enzyme which hydrolysis acetyl-CoA to acetate, but this strain was fully virulent in a mouse model of disseminated candidiasis.

7. In silico reconstruction of the sterol biosynthetic pathway and farnesol biosynthesis in *C. albicans*

The sterol biosynthesis pathway generates compounds necessary for the maintenance of cellular structure. These sterols, such as ergosterol in fungi and cholesterol in mammalian cells, are

essential for membrane integrity and permeability. The sterol biosynthesis pathway is targeted by several antifungal agents. In addition, sterol biosynthesis is potentially involved in the quorum sensing of dimorphic fungi because this pathway generates farnesol, the most well-characterized quorum sensing molecule involved in *C. albicans* morphogenesis.

Adopting the same approach we used for the central carbon metabolism, we have reconstructed, *in silico*, the *C. albicans* sterol biosynthetic pathway (Fig. 4). The first step in sterol biosynthesis is the condensation of acetyl-CoA and acetoacetyl-CoA into 3-hydroxy-3-methylglutaryl-CoA by hydroxymethylglutaryl-CoA synthase. Seven sequential reactions are then involved in the synthesis of farnesyl pyrophosphate, which is the precursor for the biosynthesis of farnesol. Surprisingly, the enzyme that synthesizes farnesol has not been fully described, but Hornby et al. (2003) claimed to have extracted an enzyme from *C. albicans* possessing this activity. Fungal and mammalian cells share a common sterol biosynthetic pathway from acetyl-CoA through farnesyl pyrophosphate to zymosterol (Fig. 4). From this point there are different pathways in fungi and mammals. 24-C-methyltransferase converts zymosterol into fecosterol leading to the biosynthesis of ergosterol in fungi, whereas 24-dehydrocholesterol reductase directs sterol biosynthesis into the production of cholesterol which serves as a precursor for steroid hormones (e.g. androgens and estrogens) in mammalian cells. In general, the biosynthesis of ergosterol is conserved between *C. albicans* and *S. cerevisiae*.

Interestingly, the presence of estrogens has been reported to have a profound effect on the morphogenesis of dimorphic fungi (Cheng et al., 2006; White and Larsen, 1997). Estrogen may contribute to the gender bias in human infection by another dimorphic and pathogenic fungus, *P. brasiliensis*, for which, contrary to *C. albicans*, the yeast form is the pathogenic state. In healthy individuals, *P. brasiliensis* infects mainly adult males (Borges-Walmsley et al., 2002). Estrogen has been shown to suppress the hyphal-to-yeast transition in *P. brasiliensis* and, thus, it has been proposed that adult women are less prone to infection by *P. brasiliensis* due to higher levels of estrogen in their bodies compared to males (Borges-Walmsley et al., 2002).

On the other hand, healthy women often develop recurrent vaginal candidosis, especially during pregnancy (Tarry et al., 2005). Clinical studies have shown that women are more likely to have symptomatic *Candida* infections when estrogen levels are high (Tarry et al., 2005), and estrogen has been shown to promote germ tube formation in *C. albicans* (Cheng et al., 2006; White and Larsen, 1997) which is the first step in the yeast-to-hypha transition (Whiteway and Bachewich, 2007). Low estrogen levels may be one reason why healthy adult males do not commonly develop candidosis.

Therefore estrogen, which seems to promote *Candida* infection in women, protects them from *P. brasiliensis* infection because this hormone induces the filamentous growth of both fungi. What is intriguing from the metabolic point of view is the fact that estrogens are synthesized by mammalian cells via the sterol biosynthetic pathway, which is used by fungi to synthesize farnesol (Nickerson et al., 2006). Thus, the sterol pathway appears to have an important role in both fungal morphogenesis and in pathogenesis.

8. The integration of quorum sensing and central carbon metabolism in *C. albicans*

QSMs are products of the central carbon metabolism of *C. albicans*. They function as communication signals and coordinately regulate the growth of *C. albicans* cells, in response to cell density. Their production, however, may simply reflect the flux within the central carbon metabolism pathways and their accumulation and




Fig. 4. The *in silico* reconstruction of the sterol pathway of *C. albicans* and mammalian cells. Farnesol, which suppress germ tube formation in *C. albicans*, is produced via the sterol pathway, but some enzymes involved in its biosynthesis are still to be identified. Mammals use the same pathway to synthesize estrogens, which promote germ tube formation in *C. albicans*.

signaling may provide feedback to these pathways to optimise growth under the particular environmental conditions. For example, if carbon and nitrogen sources are available to the cells this may stimulate flux through the pathways for the biosynthesis of QSMs such as farnesol that will repress the filamentous growth of *C. albicans* and, if carbon and/or nitrogen sources become scarce, flux through other pathways involved in the biosynthesis of QSMs that accelerate hypha formation, such as tyrosol, may be activated.

The metabolic hubs of central carbon metabolism are metabolites that take part in more than twenty different pathways and include pyruvate, NADH, NADPH, and ATP (Villas-Bôas et al., 2007). Thus, central carbon metabolism is highly interconnected and forms a metabolic network that responds very quickly to any environmental change. This is the part of cell metabolism most likely to include metabolic reactions involved in cell responses to QSMs in the environment. Indeed, several observations support the idea that central carbon metabolism provides the carbon sources for the synthesis of QSMs. As discussed above, farnesol, tyrosol, tryptophol, and phenylethyl alcohol are derived from glycolytic products. Aromatic QSMs are derived from phosphoenolpyruvate and farnesol is derived from acetyl-CoA. Therefore, if these molecules are synthesized as part of central carbon metabolism, their concentrations are likely to affect the flux distribution in these primary metabolic pathways by suppressing or inducing specific metabolic reactions. In addition, N-acyl homoserine lactones, which are common QSMs produced by many bacteria (e.g. *Pseudomonas corrugata*, *Erwinia carotovora* and others (Dong et al., 2001; Licciardello et al., 2007; Schaefer et al., 1996)), are synthesized directly from methionine, which is derived from oxaloacetic acid in the TCA cycle, and are thus also part of central carbon metabolism. **Moreover, the production of QSMs seems to be affected by the availability of their precursors. For instance, the presence of aromatic amino acids increases the level of tyrosol, tryptophol, and phenylethyl alcohol in *C. albicans* (Ghosh et al., 2008).**

Despite the obvious connection between quorum sensing and central carbon metabolism, the role of primary metabolism in

the mechanism of quorum sensing has not been investigated. We hypothesize that once extracellular QSMs reach a certain concentration, their passive diffusion across the cell membrane could affect the expression of specific genes or the activity of specific enzymes involved in morphogenesis. Indeed, fluorescently labeled farnesol analogs supplied to *C. albicans* cultures have been found in the cytoplasm of *C. albicans* (Shchepin et al., 2003). Thus, QSMs could rapidly up- and down-regulate particular metabolic pathways (e.g. aromatic amino acid biosynthesis and sterol biosynthesis), redistributing the metabolic flux of carbon through central carbon metabolism and inducing the morphological switch by altering the biosynthesis of cell wall components. Several transcriptomics studies have shown that the metabolic pathways from central carbon metabolism are highly affected by the presence of quorum sensing molecules (Table 3). Therefore, the study on the effect of QSMs on *C. albicans* morphogenesis in the context of central carbon metabolism has the potential to unravel important metabolic mechanisms underlying the morphogenetic process, and thus, should be pursued further.

9. Final remarks

The ability to study the behaviour of biological systems *in vivo* under different environmental conditions has increased with recent developments in genomics and post-genomic tools. Gene knockout experiments can demonstrate the involvement of individual genes in biological processes, but often it is unclear how the genes mediate their effects. Proteomic studies can confirm that changes in transcription associated with gene disruption, or environmental changes, result in altered protein expression. The integration of gene knockout, transcriptomic and proteomic studies has elegantly elucidated signaling pathways involved in *C. albicans* morphogenesis. This still, however, only gives a linear view of morphogenesis. This review has used the *C. albicans* genome data to

reconstruct the network of metabolic pathways involved in central carbon, sterol, and QSM metabolism. This will provide a framework which can guide the interpretation of metabolomic data and thus generate the fluxome and a more *three-dimensional* appreciation of how QSMs and environmental cues mediate their effects on morphology.

Morphogenesis is considered to be an important virulence factor for *C. albicans* and other dimorphic fungi. Central carbon metabolism and sterol biosynthesis not only supply carbon and lipid sources to generate the building blocks of new cellular structures in response to morphogenesis, but also supply precursors for the biosynthesis of the quorum sensing molecules involved in cell-cell communication and the coordinated dimorphic transition in a population of *C. albicans* cells. Together, the pre-existing genomic sequences and the *in silico* reconstruction of *C. albicans* biochemical pathways should enable a powerful systems biology study of this fungus that would be the cornerstone to assist in the elucidation of the metabolic mechanisms responsible for *C. albicans* morphogenesis and virulence.

References

- Alem, M.A.S., Oteef, M.D.Y., Flowers, T.H., Douglas, L.J., 2006. Production of tyrosol by *Candida albicans* biofilms and its role in quorum sensing and biofilm development. *Eukaryotic Cell* 5, 1770–1779.
- Alonso-Monge, R., Navarro-García, F., Román, E., Negredo, A.I., Eisman, B., Nombela, C., Pla, J., 2003. The Hog1 mitogen-activated protein kinase is essential in the oxidative stress response and chlamydospore formation in *Candida albicans*. *Eukaryotic Cell* 2, 351–361.
- Alonso-Monge, R., Carvahlo, S., Nombela, C., Rial, E., Pla, J., 2009a. The Hog1 MAP kinase controls respiratory metabolism in the fungal pathogen *Candida albicans*. *Microbiology* 155, 413–423.
- Alonso-Monge, R., Román, E., Arana, D.M., Pla, J., Nombela, C., 2009b. Fungi sensing environmental stress. *Clinical Microbiology and Infection* 15, 17–19.
- Alvarez, F.J., Konopka, J.B., 2007. Identification of an N-acetylglucosamine transporter that mediates hyphal induction in *Candida albicans*. *Molecular Biology of the Cell* 18, 965–975.
- Arai, T., Mikami, Y., Yokoyama, K., 1977. Phagocytosis of *Candida albicans* by rabbit alveolar macrophages and guinea pig neutrophils. *Sabouraudia Journal of Medical and Veterinary Mycology* 15, 171–177.
- Arana, D.M., Nombela, C., Alonso-Monge, R., Pla, J., 2005. The Pbs2 MAP kinase is essential for the oxidative-stress response in the fungal pathogen *Candida albicans*. *Microbiology* 151, 1033–1049.
- Askew, C., Sellam, A., Epp, E., Hogue, H., Mullick, A., Nantel, A., Whiteway, M., 2009. Transcriptional regulation of carbohydrate metabolism in the human pathogen *Candida albicans*. *PLoS Pathogens* 5.
- Asleson, C.M., Bensen, E.S., Gale, C.A., Melms, A.S., Kurischko, C., Berman, J., 2001. *Candida albicans* INT1-induced filamentation in *Saccharomyces cerevisiae* depends on Sla2p. *Molecular and Cellular Biology* 21, 1272–1284.
- Bachewich, C., Whiteway, M., 2005. Cyclin Cln3p links G1 progression to hyphal and pseudohyphal development in *Candida albicans*. *Eukaryotic Cell* 4, 95–102.
- Bahn, Y.S., Mühlischlegel, F.A., 2006. CO₂ sensing in fungi and beyond. *Current Opinion in Microbiology* 9, 572–578.
- Bahn, Y.S., Sundstrom, P., 2001. CAP1, an adenylate cyclase-associated protein gene, regulates bud-hypha transitions, filamentous growth, and cyclic AMP levels and is required for virulence of *Candida albicans*. *Journal of Bacteriology* 183, 3211–3223.
- Bahn, Y.S., Staab, J., Sundstrom, P., 2003. Increased high-affinity phosphodiesterase PDE2 gene expression in germ tubes counteracts CAP1-dependent synthesis of cyclic AMP, limits hypha production and promotes virulence of *Candida albicans*. *Molecular Microbiology* 50, 391–409.
- Bahn, Y.S., Xue, C., Idnurm, A., Rutherford, J.C., Heitman, J., Cardenas, M.E., 2007. Sensing the environment: lessons from fungi. *Nature Reviews Microbiology* 5, 57–69.
- Bainton, N.J., Bycroft, B.W., Chhabra, S.R., Stead, P., Gledhill, L., Hill, P.J., Rees, C.E.D., Winson, M.K., Salmond, G.P.C., Stewart, G.S.A.B., Williams, P., 1992. A general role for the lux autoinducer in bacterial cell signalling: control of antibiotic biosynthesis in *Erwinia*. *Gene* 116, 87–91.
- Banerjee, M., Thompson, D.S., Lazzell, A., Carlisle, P.L., Pierce, C., Monteagudo, C., López-Ribot, J.L., Kadosh, D., 2008. UME6, a novel filament-specific regulator of *Candida albicans* hyphal extension and virulence. *Molecular Biology of the Cell* 19, 1354–1365.
- Barelle, C.J., Priest, C.L., MacCallum, D.M., Gow, N.A.R., Odds, F.C., Brown, A.J.P., 2006. Niche-specific regulation of central metabolic pathways in a fungal pathogen. *Cellular Microbiology* 8, 961–971.
- Basiliana, M., Blyth, J., Arkowitz, R.A., 2003. Cdc24, the GDP-GTP exchange factor for Cdc42, is required for invasive hyphal growth of *Candida albicans*. *Eukaryotic Cell* 2, 9–18.
- Bastidas, R.J., Heitman, J., 2009. Trimorphic stepping stones pave the way to fungal virulence. *Proceedings of the National Academy of Sciences of the United States of America* 106, 351–352.
- Bendel, C.M., Hess, D.J., Garni, R.M., Henry-Stanley, M., Wells, C.L., 2003. Comparative virulence of *Candida albicans* yeast and filamentous forms in orally and intravenously inoculated mice. *Critical Care Medicine* 31, 501–507.
- Bensen, E.S., Martin, S.J., Li, M., Berman, J., Davis, D.A., 2004. Transcriptional profiling in *Candida albicans* reveals new adaptive responses to extracellular pH and functions for Rim101p. *Molecular Microbiology* 54, 1335–1351.
- Bensen, E.S., Clemente-Blanco, A., Finley, K.R., Correa-Bordes, J., Berman, J., 2005. The mitotic cyclins Clb2p and Clb4p affect morphogenesis in *Candida albicans*. *Molecular Biology of the Cell* 16, 3387–3400.
- Biswas, K., Morschhäuser, J., 2005. The Mep2p ammonium permease controls nitrogen starvation-induced filamentous growth in *Candida albicans*. *Molecular Microbiology* 56, 649–669.
- Biswas, S., Roy, M., Datta, A., 2003. N-acetylglucosamine-inducible CaGAP1 encodes a general amino acid permease which co-ordinates external nitrogen source response and morphogenesis in *Candida albicans*. *Microbiology* 149, 2597–2608.
- Biswas, S., Van Dijk, P., Datta, A., 2007. Environmental sensing and signal transduction pathways regulating morphopathogenic determinants of *Candida albicans*. *Microbiology and Molecular Biology Reviews* 71, 348–376.
- Borges-Walmsley, M.L., Chen, D., Shu, X., Walmsley, A.R., 2002. The pathobiology of *Paracoccidioides brasiliensis*. *Trends in Microbiology* 10, 80–87.
- Braun, B.R., Johnson, A.D., 1997. Control of filament formation in *Candida albicans* by the transcriptional repressor TUP1. *Science* 277, 105–109.
- Braun, B.R., Kadosh, D., Johnson, A.D., 2001. NRG1, a repressor of filamentous growth in *C. albicans*, is down-regulated during filament induction. *EMBO Journal* 20, 4753–4761.
- Brown, A.J.P., Gow, N.A.R., 1999. Regulatory networks controlling *Candida albicans* morphogenesis. *Trends in Microbiology* 7, 333–338.
- Brown Jr., D.H., Giusani, A.D., Chen, X., Kumamoto, C.A., 1999. Filamentous growth of *Candida albicans* in response to physical environmental cues and its regulation by the unique CZF1 gene. *Molecular Microbiology* 34, 651–662.
- Brown, V., Sexton, J.A., Johnston, M., 2006. A glucose sensor in *Candida albicans*. *Eukaryotic Cell* 5, 1726–1737.
- Brown, V., Sabina, J., Johnston, M., 2009. Specialized sugar sensing in diverse fungi. *Current Biology* 19, 436–441.
- Cabezón, V., Llama-Palacios, A., Nombela, C., Monteoliva, L., Gil, C., 2009. Analysis of *Candida albicans* plasma membrane proteome. *Proteomics* 9, 4770–4786.
- Calera, J.A., Calderone, R., 1999. Histidine kinase, two-component signal transduction proteins of *Candida albicans* and the pathogenesis of candidosis. *Mycoses* 42 (1), 49–53 (Suppl.).
- Calera, J.A., Zhao, X.J., Calderone, R., 2000. Defective hyphal development and avirulence caused by a deletion of the SSK1 response regulator gene in *Candida albicans*. *Infection and Immunity* 68, 518–525.
- Cao, Y.Y., Cao, Y.B., Xu, Z., Ying, K., Li, Y., Xie, Y., Zhu, Z.Y., Chen, W.S., Jiang, Y.Y., 2005. CDNA microarray analysis of differential gene expression in *Candida albicans* biofilm exposed to farnesol. *Antimicrobial Agents and Chemotherapy* 49, 584–589.
- Carlisle, P.L., Banerjee, M., Lazzell, A., Monteagudo, C., López-Ribot, J.L., Kadosh, D., 2009. Expression levels of a filament-specific transcriptional regulator are sufficient to determine *Candida albicans* morphology and virulence. *Proceedings of the National Academy of Sciences of the United States of America* 106, 599–604.
- Carman, A.J., Vylkova, S., Lorenz, M.C., 2008. Role of acetyl coenzyme A synthesis and breakdown in alternative carbon source utilization in *Candida albicans*. *Eukaryotic Cell* 7, 1733–1741.
- Cassola, A., Parrot, M., Silberstein, S., Magee, B.B., Passeron, S., Giasson, L., Cantore, M.L., 2004. *Candida albicans* lacking the gene encoding the regulatory subunit of protein kinase A displays a defect in hyphal formation and an altered localization of the catalytic subunit. *Eukaryotic Cell* 3, 190–199.
- Cheetham, J., Smith, D.A., Dantas, A.D.S., Doris, K.S., Patterson, M.J., Bruce, C.R., Quinn, J., 2007. A single MAPKKK regulates the Hog1 MAPK pathway in the pathogenic fungus *Candida albicans*. *Molecular Biology of the Cell* 18, 4603–4614.
- Chen, H., Fink, G.R., 2006. Feedback control of morphogenesis in fungi by aromatic alcohols. *Genes & Development* 20, 1150–1161.
- Chen, D., Janganan, T.K., Chen, G., Marques, E.R., Kress, M.R., Goldman, G.H., Walmsley, A.R., Borges-Walmsley, M.L., 2007. The cAMP pathway is important for controlling the morphological switch to the pathogenic yeast form of *Paracoccidioides brasiliensis*. *Molecular Microbiology* 65, 761–779.
- Chen, J., Zhou, S., Wang, Q., Chen, X., Pan, T., Liu, H., 2000. Crk1, a novel Cdc2-related protein kinase, is required for hyphal development and virulence in *Candida albicans*. *Molecular and Cellular Biology* 20, 8696–8708.
- Chen, H., Fujita, M., Feng, Q., Clardy, J., Fink, G.R., 2004. Tyrosol is a quorum-sensing molecule in *Candida albicans*. *Proceedings of the National Academy of Sciences of the United States of America* 101, 5048–5052.
- Cheng, G., Yeater, K.M., Hoyer, L.L., 2006. Cellular and molecular biology of *Candida albicans* estrogen response. *Eukaryotic Cell* 5, 180–191.
- Cloutier, M., Castilla, R., Bolduc, N., Zelada, A., Martineau, P., Bouillon, M., Magee, B.B., Passeron, S., Giasson, L., Cantore, M.L., 2003. The two isoforms of the cAMP-dependent protein kinase catalytic subunit are involved in the control of dimorphism in the human fungal pathogen *Candida albicans*. *Fungal Genetics and Biology* 38, 133–141.

- Court, H., Sudbery, P., 2007. Regulation of Cdc42 GTPase activity in the formation of hyphae in *Candida albicans*. *Molecular Biology of the Cell* 18, 265–281.
- Cremer, J., Vatou, V., Braveny, I., 1999. 2,4-(Hydroxyphenyl)-ethanol, an antioxidant agent produced by *Candida* spp., impairs neutrophilic yeast killing in vitro. *FEMS Microbiology Letters* 170, 319–325.
- Csank, C., Makris, C., Meloche, S., Schroppel, K., Rollinghoff, M., Dignard, D., Thomas, D.Y., Whiteway, M., 1997. Derepressed hyphal growth and reduced virulence in a VH1 family-related protein phosphatase mutant of the human pathogen *Candida albicans*. *Molecular Biology of the Cell* 8, 2539–2551.
- Csank, C., Schröppel, K., Leberer, E., Harscus, D., Mohamed, O., Meloche, S., Thomas, D.Y., Whiteway, M., 1998. Roles of the *Candida albicans* mitogen-activated protein kinase homolog, Cek1p, in hyphal development and systemic candidiasis. *Infection and Immunity* 66, 2713–2721.
- Dabrowa, N., Taxer, S.S.S., Howard, D.H., 1976. Germination of *Candida albicans* induced by proline. *Infection and Immunity* 13, 830–835.
- Davis-Hanna, A., Piispanen, A.E., Stateva, L.L., Hogan, D.A., 2008. Farnesol and dodecanol effects on the *Candida albicans* Ras1-cAMP signalling pathway and the regulation of morphogenesis. *Molecular Microbiology* 67, 47–62.
- De Bernardis, F., Mühlischlegel, F.A., Cassone, A., Fonzi, W.A., 1998. The pH of the host niche controls gene expression in and virulence of *Candida albicans*. *Infection and Immunity* 66, 3317–3325.
- Décasis, N., Savignac, K., Rouabhi, M., 2009. Farnesol promotes epithelial cell defense against *Candida albicans* through Toll-like receptor 2 expression, interleukin-6 and human β -defensin 2 production. *Cytokine* 45, 132–140.
- Derengowski, L.S., De-Souza-Silva, C., Braz, S.V., Mello-De-Sousa, T.M., Bão, S.N., Kyaw, C.M., Silva-Pereira, I., 2009. Antimicrobial effect of farnesol, a *Candida albicans* quorum sensing molecule, on Paracoccidioides brasiliensis growth and morphogenesis. *Annals of Clinical Microbiology and Antimicrobials*, 8.
- Dhillon, N.K., Sharma, S., Khuller, G.K., 2003. Signaling through protein kinases and transcriptional regulators in *Candida albicans*. *Critical Reviews in Microbiology* 29, 259–275.
- Doedt, T., Krishnamurthy, S., Bockmühl, D.P., Tebarth, B., Stempel, C., Russell, C.L., Brown, A.J.P., Ernst, J.F., 2004. APSES proteins regulate morphogenesis and metabolism in *Candida albicans*. *Molecular Biology of the Cell* 15, 3167–3180.
- Dong, Y.H., Wang, L.H., Xu, J.L., Zhang, H.B., Zhang, X.F., Zhang, L.H., 2001. Quenching quorum-sensing-dependent bacterial infection by an N-acyl homoserine lactonase. *Nature* 411, 813–817.
- Eberhard, A., 1972. Inhibition and activation of bacterial luciferase synthesis. *Journal of Bacteriology* 109, 1101–1105.
- Eberhard, A., Burlingame, A.L., Eberhard, C., Kenyon, G.L., Neilson, K.H., Oppenheimer, N.J., 1981. Structural identification of autoinducer of *Photobacterium fischeri* luciferase. *Biochemistry* 20, 2444–2449.
- Eisman, B., Alonso-Monge, R., Román, E., Arana, D., Nombela, C., Pla, J., 2006. The Cek1 and Hog1 mitogen-activated protein kinases play complementary roles in cell wall biogenesis and chlamydospore formation in the fungal pathogen *Candida albicans*. *Eukaryotic Cell* 5, 347–358.
- Enjalbert, B., Whiteway, M., 2005. Release from quorum-sensing molecules triggers hyphal formation during *Candida albicans* resumption of growth. *Eukaryotic Cell* 4, 1203–1210.
- Enjalbert, B., Smith, D.A., Cornell, M.J., Alam, I., Nicholls, S., Brown, A.J.P., Quinn, J., 2006. Role of the Hog1 stress-activated protein kinase in the global transcriptional response to stress in the fungal pathogen *Candida albicans*. *Molecular Biology of the Cell* 17, 1018–1032.
- Enloe, B., Diamond, A., Mitchell, A.P., 2000. A single-transformation gene function test in diploid *Candida albicans*. *Journal of Bacteriology* 182, 5730–5736.
- Ernst, J.F., 2000. Transcription factors in *Candida albicans*-environmental control of morphogenesis. *Microbiology* 146, 1763–1774.
- Ernandes, J.R., De Meirsmen, C., Rolland, F., Winderickx, J., De Winde, J., Brandão, R.L., Thevelein, J.M., 1998. During the initiation of fermentation overexpression of hexokinase PII in yeast transiently causes a similar deregulation of glycolysis as deletion of Tps1. *Yeast* 14, 255–269.
- Fairn, G.D., MacDonald, K., McMaster, C.R., 2007. A chemogenomic screen in *Saccharomyces cerevisiae* uncovers a primary role for the mitochondria in farnesol toxicity and its regulation by the Pkc1 pathway. *Journal of Biological Chemistry* 282, 4868–4874.
- Felix, E., 1907. Über die Bedingungen der Fuselölbildung und über ihren Zusammenhang mit dem Eiweißaufbau der Hefe. *Berichte der Deutschen Chemischen Gesellschaft* 40, 1027–1047.
- Feng, Q., Summers, E., Guo, B., Fink, G., 1999. Ras signaling is required for serum-induced hyphal differentiation in *Candida albicans*. *Journal of Bacteriology* 181, 6339–6346.
- Fernández-Arenas, E., Cabezón, V., Bermejo, C., Arroyo, J., Nombela, C., Diez-Orejas, R., Gil, C., 2007. Integrated proteomics and genomics strategies bring new insight into *Candida albicans* response upon macrophage interaction. *Molecular and Cellular Proteomics* 6, 460–478.
- Fradin, C., De Groot, P., MacCallum, D., Schaller, M., Klis, F., Odds, F.C., Hube, B., 2005. Granulocytes govern the transcriptional response, morphology and proliferation of *Candida albicans* in human blood. *Molecular Microbiology* 56, 397–415.
- Fu, Y., Ibrahim, A.S., Sheppard, D.C., Chen, Y.C., French, S.W., Cutler, J.E., Filler, S.G., Edwards Jr., J.E., 2002. *Candida albicans* Als1p: an adhesin that is a downstream effector of the EFG1 filamentation pathway. *Molecular Microbiology* 44, 61–72.
- Fuqua, W.C., Winans, S.C., Greenberg, E.P., 1994. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *Journal of Bacteriology* 176, 269–275.
- Gale, C.A., Bendel, C.M., McClellan, M., Hauser, M., Becker, J.M., Berman, J., Hostetter, M.K., 1998. Linkage of adhesion, filamentous growth, and virulence in *Candida albicans* to a single gene, INT1. *Science* 279, 1355–1358.
- García-Sánchez, S., Aubert, S., Iraqui, I., Janbon, G., Ghigo, J.M., D'Enfert, C., 2004. *Candida albicans* biofilms: a developmental state associated with specific and stable gene expression patterns. *Eukaryotic Cell* 3, 536–545.
- García-Sánchez, S., Mavor, A.L., Russell, C.L., Argimon, S., Dennison, P., Enjalbert, B., Brown, A.J.P., 2005. Global roles of Ssn6 in tup1- and nrg1-dependent gene regulation in the fungal pathogen, *Candida albicans*. *Molecular Biology of the Cell* 16, 2913–2925.
- Channoum, M.A., Spellberg, B., Saporito-Irwin, S.M., Fonzi, W.A., 1995. Reduced virulence of *Candida albicans* PHR1 mutants. *Infection and Immunity* 63, 4528–4530.
- Ghosh, S., Howe, N., Volk, K., Tati, S., Nickerson, K.W., Petro, T.M., 2010. *Candida albicans* cell wall components and farnesol stimulate the expression of both inflammatory and regulatory cytokines in the murine RAW264.7 macrophage cell line. *FEMS Immunology and Medical Microbiology* 60, 63–73.
- Ghosh, S., Kebara, B.W., Atkin, A.L., Nickerson, K.W., 2008. Regulation of aromatic alcohol production in *Candida albicans*. *Applied and Environmental Microbiology* 74, 7211–7218.
- Ghosh, S., Navarathna, D.H.M.L.P., Roberts, D.D., Cooper, J.T., Atkin, A.L., Petro, T.M., Nickerson, K.W., 2009. Arginine-induced germ tube formation in *Candida albicans* is essential for escape from murine macrophage line RAW 264.7. *Infection and Immunity* 77, 1596–1605.
- Gow, N.A.R., 2009. Fungal morphogenesis: some like it hot. *Current Biology* 19, 621–629.
- Gow, N.A.R., Brown, A.J.P., Odds, F.C., 2002. Fungal morphogenesis and host invasion. *Current Opinion in Microbiology* 5, 366–371.
- Goyard, S., Knechtle, P., Chauvel, M., Mallet, A., Prevost, M.-C., Proux, C., Coppee, J.-Y., Schwartz, P., Dromer, F., Park, H., Filler, S.G., Janbon, G., d'Enfert, C., 2008. The Yak1 kinase is involved in the initiation and maintenance of hyphal growth in *Candida albicans*. *Molecular Biology of the Cell* 19, 2251–2266.
- Gray, K.M., Garey, J.R., 2001. The evolution of bacterial LuxI and LuxR quorum sensing regulators. *Microbiology* 147, 2379–2387.
- Gray, K.M., Passador, L., Iglewski, B.H., Greenberg, E.P., 1994. Interchangeability and specificity of components from the quorum-sensing regulatory systems of *Vibrio fischeri* and *Pseudomonas aeruginosa*. *Journal of Bacteriology* 176, 3076–3080.
- Grubb, S.E.W., Murdoch, C., Sudbery, P.E., Saville, S.P., Lopez-Ribot, J.L., Thornhill, M.B., 2009. Adhesion of *Candida albicans* to endothelial cells under physiological conditions of flow. *Infection and Immunity* 77, 3872–3878.
- Hairfield, M.L., Westwater, C., Dolan, J.W., 2002. Phosphatidylinositol-4-phosphate 5-kinase activity is stimulated during temperature-induced morphogenesis in *Candida albicans*. *Microbiology* 148, 1737–1746.
- Harcus, D., Nantel, A., Marciel, A., Rigby, T., Whiteway, M., 2004. Transcription profiling of cyclic AMP signaling in *Candida albicans*. *Molecular Biology of the Cell* 15, 4490–4499.
- Hazelwood, L.A., Daran, J.M., Van Maris, A.J.A., Pronk, J.T., Dickinson, J.R., 2008. The Ehrlich pathway for fusel alcohol production: a century of research on *Saccharomyces cerevisiae* metabolism. *Applied and Environmental Microbiology* 74, 2259–2266.
- Henriques, M., Martins, M., Azeredo, J., Oliveira, R., 2007. Effect of farnesol on *Candida dubliniensis* morphogenesis. *Letters in Applied Microbiology* 44, 199–205.
- Hense, B.A., Kuttler, C., Müller, J., Rothballer, M., Hartmann, A., Kreft, J.U., 2007. Does efficiency sensing unify diffusion and quorum sensing? *Nature Reviews Microbiology* 5, 230–239.
- Hernández, R., Nombela, C., Diez-Orejas, R., Gil, C., 2004. Two-dimensional reference map of *Candida albicans* hyphal forms. *Proteomics* 4, 374–382.
- Hilmioglu, S., Ilkit, M., Badak, Z., 2007. Comparison of 12 liquid media for germ tube production of *Candida albicans* and *C. tropicalis*. *Mycoses* 50, 282–285.
- Hogan, D.A., 2006a. Quorum sensing: alcohols in a social situation. *Current Biology*, 16.
- Hogan, D.A., 2006b. Talking to themselves: autoregulation and quorum sensing in fungi. *Eukaryotic Cell* 5, 613–619.
- Hogan, D.A., Sundstrom, P., 2009. The Ras/cAMP/PKA signaling pathway and virulence in *Candida albicans*. *Future Microbiology* 4, 1263–1270.
- Hogan, D.A., Vik, A., Kolter, R., 2004. A *Pseudomonas aeruginosa* quorum-sensing molecule influences *Candida albicans* morphology. *Molecular Microbiology* 54, 1212–1223.
- Holmes, A.R., Shepherd, M.G., 1987. Proline-induced germ-tube formation in *Candida albicans*: role of proline uptake and nitrogen metabolism. *Journal of General Microbiology* 133, 3219–3228.
- Holmes, A.R., Cannon, R.D., Shepherd, M.G., 1991. Effect of calcium ion uptake on *Candida albicans* morphology. *FEMS Microbiology Letters* 77, 187–193.
- Hornby, J.M., Kebara, B.W., Nickerson, K.W., 2003. Farnesol biosynthesis in *Candida albicans*: Cellular response to sterol inhibition by zaragozic acid B. *Antimicrobial Agents and Chemotherapy* 47, 2366–2369.
- Hornby, J.M., Nickerson, K.W., 2004. Enhanced production of farnesol by *Candida albicans* treated with four azoles. *Antimicrobial Agents and Chemotherapy* 48, 2305–2307.
- Hornby, J.M., Jensen, E.C., Lisec, A.D., Tasto, J.J., Jahnke, B., Shoemaker, R., Dussault, P., Nickerson, K.W., 2001. Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. *Applied and Environmental Microbiology* 67, 2982–2992.
- Hrmova, M., Drobnica, L., 1981. Induction of mycelial type of development in *Candida albicans* by low glucose concentration. *Mycopathologia* 76, 83–96.

- Hromatka, B.S., Noble, S.M., Johnson, A.D., 2005. Transcriptional response of *Candida albicans* to nitric oxide and the role of the YHB1 gene in nitrosative stress and virulence. *Molecular Biology of the Cell* 16, 4814–4826.
- Hudson, D.A., Sciascia, Q.L., Sanders, R.J., Norris, G.E., Edwards, P.J.B., Sullivan, P.A., Farley, P.C., 2004. Identification of the dialysable serum inducer of germ-tube formation in *Candida albicans*. *Microbiology* 150, 3041–3049.
- Huh, W.K., Kim, S.T., Kim, H., Jeong, G., Kang, S.O., 2001. Deficiency of D-erythroascorbic acid attenuates hyphal growth and virulence of *Candida albicans*. *Infection and Immunity* 69, 3939–3946.
- Hwang, C.S., Oh, J.H., Huh, W.K., Yim, H.S., Kang, S.O., 2003. Ssn6, an important factor of morphological conversion and virulence in *Candida albicans*. *Molecular Microbiology* 47, 1029–1043.
- Johnson, D.I., 1999. Cdc42: an essential Rho-type GTPase controlling eukaryotic cell polarity. *Microbiology and Molecular Biology Reviews* 63, 54–105.
- José, C.S., Monge, R.A., Pérez-Díaz, R., Pla, J., Nombela, C., 1996. The mitogen-activated protein kinase homolog HOG1 gene controls glycerol accumulation in the pathogenic fungus *Candida albicans*. *Journal of Bacteriology* 178, 5850–5852.
- Kadosh, D., Johnson, A.D., 2001. Rfg1, a protein related to the *Saccharomyces cerevisiae* hypoxic regulator Rox1, controls filamentous growth and virulence in *Candida albicans*. *Molecular and Cellular Biology* 21, 2496–2505.
- Karababa, M., Coste, A.T., Rognon, B., Bille, J., Sanglard, D., 2004. Comparison of gene expression profiles of *Candida albicans* azole-resistant clinical isolates and laboratory strains exposed to drugs inducing multidrug transporters. *Antimicrobial Agents and Chemotherapy* 48, 3064–3079.
- Kebaara, B.W., Langford, M.L., Navarathna, D.H.M.L.P., Dumitru, R., Nickerson, K.W., Atkin, A.L., 2008. *Candida albicans* Tup1 is involved in farnesol-mediated inhibition of filamentous-growth induction. *Eukaryotic Cell* 7, 980–987.
- Khalaf, R.A., Zitomer, R.S., 2001. The DNA binding protein Rfg1 is a repressor of filamentation in *Candida albicans*. *Genetics* 157, 1503–1512.
- Klengel, T., Liang, W.-J., Chaloupka, J., Ruoff, C., Schröpel, K., Naglik, J.R., Eckert, S.E., Mogensen, E.G., Haynes, K., Tuite, M.F., Levin, L.R., Buck, J., Mülschlegel, F.A., 2005. Fungal adenyl cyclase integrates CO₂ sensing with cAMP signaling and virulence. *Current Biology* 15, 2021–2026.
- Kruppa, M., Krom, B.P., Chauhan, N., Bambach, A.V., Cihlar, R.L., Calderone, R.A., 2004. The two-component signal transduction protein Chk1p regulates quorum sensing in *Candida albicans*. *Eukaryotic Cell* 3, 1062–1065.
- Kügler, S., Sebghati, T.S., Eissenberg, L.G., Goldman, W.E., 2000. Phenotypic variation and intracellular parasitism by *Histoplasma capsulatum*. *Proceedings of the National Academy of Sciences of the United States of America* 97, 8794–8798.
- Kusch, H., Engelmann, S., Bode, R., Albrecht, D., Morschhäuser, J., Hecker, M., 2008. A proteomic view of *Candida albicans* yeast cell metabolism in exponential and stationary growth phases. *International Journal of Medical Microbiology* 298, 291–318.
- Lan, C.Y., Rodarte, G., Murillo, L.A., Jones, T., Davis, R.W., Dungan, J., Newport, G., Agabian, N., 2004. Regulatory networks affected by iron availability in *Candida albicans*. *Molecular Microbiology* 53, 1451–1469.
- Lane, S., Birse, C., Zhou, S., Matson, R., Liu, H., 2001. DNA array studies demonstrate convergent regulation of virulence factors by Cph1, Cph2, and Efg1 in *Candida albicans*. *Journal of Biological Chemistry* 276, 48988–48996.
- Langford, M.L., Hasim, S., Nickerson, K.W., Atkin, A.L., 2010. Activity and toxicity of farnesol towards *Candida albicans* are dependent on growth conditions. *Antimicrobial Agents and Chemotherapy* 54, 940–942.
- Lazo, Chapa Y., Bates, S., Sudbery, P., 2005. The G1 cyclin Cln3 regulates morphogenesis in *Candida albicans*. *Eukaryotic Cell* 4, 90–94.
- Leberer, E., Harcus, D., Broadbent, I.D., Clark, K.L., Dignard, D., Ziegelbauer, K., Schmidt, A., Gow, N.A.R., Brown, A.J.P., Thomas, D.Y., 1996. Signal transduction through homologs of the Ste20p and Ste7p protein kinases can trigger hyphal formation in the pathogenic fungus *Candida albicans*. *Proceedings of the National Academy of Sciences of the United States of America* 93, 13217–13222.
- Leberer, E., Harcus, D., Dignard, D., Johnson, L., Ushinsky, S., Thomas, D.Y., Schröpel, K., 2001. Ras links cellular morphogenesis to virulence by regulation of the MAP kinase and cAMP signalling pathways in the pathogenic fungus *Candida albicans*. *Molecular Microbiology* 42, 673–687.
- Lee, H., Mitchell, D.R.S., 1979. Commitment to germ tube or bud formation during release from stationary phase in *Candida albicans*. *Experimental Cell Research* 120, 167–179.
- Lee, K.L., Buckley, H.R., Campbell, C.C., 1975. An amino acid liquid synthetic medium for the development of mycelial and yeast forms of *Candida albicans*. *Sabouraudia Journal of Medical and Veterinary Mycology* 13, 148–153.
- Lee, K.H., Shin, W.S., Kim, D., Koh, C.M., 1999. The presumptive identification of *Candida albicans* with germ tube induced by high temperature. *Yonsei Medical Journal* 40, 420–424.
- Leng, P., Lee, P.R., Wu, H., Brown, A.J.P., 2001. Efg1, a morphogenetic regulator in *Candida albicans*, is a sequence-specific DNA binding protein. *Journal of Bacteriology* 183, 4090–4093.
- Lengeler, K.B., Davidson, R.C., D'Souza, C., Harashima, T., Shen, W.C., Wang, P., Pan, X., Waugh, M., Heitman, J., 2000. Signal transduction cascades regulating fungal development and virulence. *Microbiology and Molecular Biology Reviews* 64, 746–785.
- Leroy, O., Gangneux, J.P., Montravers, P., Mira, J.P., Gouin, F., Sollet, J.P., Carlet, J., Reynes, J., Rosenheim, M., Regnier, B., Lortholary, O., 2009. Epidemiology, management, and risk factors for death of invasive *Candida* infections in critical care: a multicenter, prospective, observational study in France (2005–2006). *Critical Care Medicine* 37, 1612–1618.
- Li, W., Mitchell, A.P., 1997. Proteolytic activation of rim1p, a positive regulator of yeast sporulation and invasive growth. *Genetics* 145, 63–73.
- Licciardello, G., Bertani, I., Steindler, L., Bella, P., Venturi, V., Catara, V., 2007. *Pseudomonas corrugata* contains a conserved N-acyl homoserine lactone quorum sensing system; its role in tomato pathogenicity and tobacco hypersensitivity response. *FEMS Microbiology Ecology* 61, 222–234.
- Lingappa, B.T., Prasad, M., Lingappa, Y., Hunt, D.F., Biemann, K., 1969. Phenethyl alcohol and tryptophol: antibiotics produced by the fungus *Candida albicans*. *Science* 163, 192–194.
- Liu, H., 2001. Transcriptional control of dimorphism in *Candida albicans*. *Current Opinion in Microbiology* 4, 728–735.
- Liu, H., 2002. Co-regulation of pathogenesis with dimorphism and phenotypic switching in *Candida albicans*, a commensal and a pathogen. *International Journal of Medical Microbiology* 292, 299–311.
- Liu, H., Kohler, J., Fink, G.R., 1994. Suppression of hyphal formation in *Candida albicans* by mutation of a STE12 homolog. *Science* 266, 1723–1726.
- Liu, P., Deng, B., Long, C.A., Min, X., 2009. Effect of farnesol on morphogenesis in the fungal pathogen *Penicillium expansum*. *Annals of Microbiology* 59, 33–38.
- Lo, H.J., Köhler, J.R., Didomenico, B., Loebenberg, D., Cacciapuoti, A., Fink, G.R., 1997. Nonfilamentous *C. albicans* mutants are avirulent. *Cell* 90, 939–949.
- Loeb, J.D.J., Sepúlveda-Becerra, M., Hazan, I., Liu, H., 1999. A G1 cyclin is necessary for maintenance of filamentous growth in *Candida albicans*. *Molecular and Cellular Biology* 19, 4019–4027.
- Lorenz, M.C., Fink, G.R., 2001. The glyoxylate cycle is required for fungal virulence. *Nature* 412, 83–86.
- Lorenz, M.C., Bender, J.A., Fink, G.R., 2004. Transcriptional response of *Candida albicans* upon internalization by macrophages. *Eukaryotic Cell* 3, 1076–1087.
- Maidan, M.M., De Rop, L., Serneels, J., Exler, S., Rupp, S., Tournu, H., Thevelein, J.M., Van Dijck, P., 2005a. The G protein-coupled receptor Gpr1 and the Gα protein Gpa2 act through the cAMP-protein kinase A pathway to induce morphogenesis in *Candida albicans*. *Molecular Biology of the Cell* 16, 1971–1986.
- Maidan, M.M., Thevelein, J.M., Van Dijck, P., 2005b. Carbon source induced yeast-to-hypha transition in *Candida albicans* is dependent on the presence of amino acids and on the G-protein-coupled receptor Gpr1. *Biochemical Society Transactions* 33, 291–293.
- Martins, L.F., Montero-Lomeli, M., Masuda, C.A., Fortes, F.S.A., Previato, J.O., Mendonça-Previato, L., 2008. Lithium-mediated suppression of morphogenesis and growth in *Candida albicans*. *FEMS Yeast Research* 8, 615–621.
- Mavor, A.L., Thewes, S., Hube, B., 2005. Systemic fungal infections caused by *Candida* species: epidemiology, infection process and virulence attributes. *Current Drug Targets* 6, 863–874.
- Menon, V., Li, D., Chauhan, N., Rajnarayanan, R., Dubrovskaya, A., West, A.H., Calderone, R., 2006. Functional studies of the Ssk1p response regulator protein of *Candida albicans* as determined by phenotypic analysis of receiver domain point mutants. *Molecular Microbiology* 62, 997–1013.
- Mitchell, A.P., 1998. Dimorphism and virulence in *Candida albicans*. *Current Opinion in Microbiology* 1, 687–692.
- Mock, R.C., Pollack, J.H., Hashimoto, T., 1990. Carbon dioxide induces endotrophic germ tube formation in *Candida albicans*. *Canadian Journal of Microbiology* 36, 249–253.
- Moran, C., Grussemeyer, C.A., Spalding, J.R., Benjamin Jr., D.K., Reed, S.D., 2010. Comparison of costs, length of stay, and mortality associated with *Candida glabrata* and *Candida albicans* bloodstream infections. *American Journal of Infection Control* 38, 78–80.
- Mosel, D.D., Dumitru, R., Hornby, J.M., Atkin, A.L., Nickerson, K.W., 2005. Farnesol concentrations required to block germ tube formation in *Candida albicans* in the presence and absence of serum. *Applied and Environmental Microbiology* 71, 4938–4940.
- Mühlschlegel, F.A., Fonzi, W.A., 1997. PHR2 of *Candida albicans* encodes a functional homolog of the pH-regulated gene PHR1 with an inverted pattern of pH-dependent expression. *Molecular and Cellular Biology* 17, 5960–5967.
- Murad, A.M.A., D'Enfert, C., Gaillardin, C., Tournu, H., Tekia, F., Talibi, D., Marechal, D., Marchais, V., Cottin, J., Brown, A.J.P., 2001. Transcript profiling in *Candida albicans* reveals new cellular functions for the transcriptional repressors CaTup1, CaMig1 and CaNrg1. *Molecular Microbiology* 42, 981–993.
- Nagahashi, S., Mio, T., Ono, N., Yamada-Okabe, T., Arisawa, M., Bussey, H., Yamada-Okabe, H., 1998. Isolation of CaSLN1 and CaNIK1, the genes for osmosensing histidine kinase homologues, from the pathogenic fungus *Candida albicans*. *Microbiology* 144, 425–432.
- Nantel, A., Dignard, D., Bachewich, C., Harcus, D., Marciel, A., Bouin, A.P., Sensen, C.W., Hogues, H., Van het Hoog, M., Gordon, P., Rigby, T., Benoit, F., Tessier, D.C., Thomas, D.Y., Whiteway, M., 2002. Transcript profiling of *Candida albicans* cells undergoing the yeast-to-hyphal transition. *Molecular Biology of the Cell* 13, 3452–3465.
- Navarathna, D.H.M.L.P., Hornby, J.M., Hoerrmann, N., Parkhurst, A.M., Duhamel, G.E., Nickerson, K.W., 2005. Enhanced pathogenicity of *Candida albicans* pre-treated with subinhibitory concentrations of fluconazole in a mouse model of disseminated candidiasis. *Journal of Antimicrobial Chemotherapy* 56, 1156–1159.
- Navarro-García, F., Alonso-Monge, R., Rico, H., Pla, J., Sentandreu, R., Nombela, C., 1998. A role for the MAP kinase gene MKC1 in cell wall construction and morphological transitions in *Candida albicans*. *Microbiology* 144, 411–424.
- Navarro-García, F., Sánchez, M., Nombela, C., Pla, J., 2001. Virulence genes in the pathogenic yeast *Candida albicans*. *FEMS Microbiology Reviews* 25, 245–268.
- Nealson, K.H., 1977. Autoinduction of bacterial luciferase. Occurrence, mechanism and significance. *Archives of Microbiology* 112, 73–79.
- Nemecek, J.C., Wüthrich, M., Klein, B.S., 2006. Global control of dimorphism and virulence in fungi. *Science* 312, 583–588.

- Nett, J.E., Lepak, A.J., Marchillo, K., Andes, D.R., 2009. Time course global gene expression analysis of an in vivo *Candida* biofilm. *Journal of Infectious Diseases* 200, 307–313.
- Nickerson, K.W., Atkin, A.L., Hornby, J.M., 2006. Quorum sensing in dimorphic fungi: farnesol and beyond. *Applied and Environmental Microbiology* 72, 3805–3813.
- Niimi, M., Niimi, K., Tokunaga, J., Nakayama, H., 1980. Changes in cyclic nucleotide levels and dimorphic transition in *Candida albicans*. *Journal of Bacteriology* 142, 1010–1014.
- Nobile, C.J., Solis, N., Myers, C.L., Fay, A.J., Deneault, J.S., Nantel, A., Mitchell, A.P., Filler, S.G., 2008. *Candida albicans* transcription factor Rim101 mediates pathogenic interactions through cell wall functions. *Cellular Microbiology* 10, 2180–2196.
- Odds, F.C., 1988. *Candida and Candidosis: A Review and Bibliography*, London.
- Oh, K.B., Miyazawa, H., Naito, T., Matsuoka, H., 2001. Purification and characterization of an autoregulatory substance capable of regulating the morphological transition in *Candida albicans*. *Proceedings of the National Academy of Sciences of the United States of America* 98, 4664–4668.
- Padovan, A.C.B., Chaves, G.M., Colombo, A.L., Briones, M.R.S., 2009. A novel allele of HWP1, isolated from a clinical strain of *Candida albicans* with defective hyphal growth and biofilm formation, has deletions of Gln/Pro and Ser/Thr repeats involved in cellular adhesion. *Medical Mycology* 47, 824–835.
- Payne, W.E., Kaiser, C.A., Bevis, B.J., Soderholm, J., Fu, D., Sears, I.B., Glick, B.S., 2000. Identification of YPDI, a gene of *Candida albicans* which encodes a two-component phosphohistidine intermediate protein. *Yeast* 16, 1053–1059.
- Pedreño, Y., Maicas, S., Argüelles, J.C., Sentandreu, R., Valentín, E., 2004. The ATC1 gene encodes a cell wall-linked acid trehalase required for growth on trehalose in *Candida albicans*. *Journal of Biological Chemistry* 279, 40852–40860.
- Pedreño, Y., González-Párraga, P., Martínez-Esparza, M., Sentandreu, R., Valentín, E., Argüelles, J.C., 2007. Disruption of the *Candida albicans* ATC1 gene encoding a cell-linked acid trehalase decreases hypha formation and infectivity without affecting resistance to oxidative stress. *Microbiology* 153, 1372–1381.
- Pfaller, M.A., Diekema, D.J., 2007. Epidemiology of invasive candidiasis: a persistent public health problem. *Clinical Microbiology Reviews* 20, 133–163.
- Piekarska, K., Mol, E., Van Den Berg, M., Hardy, G., Van Den Burg, J., Van Roermund, C., MacCallum, D., Odds, F., Distel, B., 2006. Peroxisomal fatty acid β -oxidation is not essential for virulence of *Candida albicans*. *Eukaryotic Cell* 5, 1847–1856.
- Piper, K.R., Beck Von Bodman, S., Farrand, S.K., 1993. Conjugation factor of *Agrobacterium tumefaciens* regulates Ti plasmid transfer by autoinduction. *Nature* 362, 448–450.
- Pitarch, A., Abian, J., Carrascal, M., Sánchez, M., Nombela, C., Gil, C., 2004. Proteomics-based identification of novel *Candida albicans* antigens for diagnosis of systemic candidiasis in patients with underlying hematological malignancies. *Proteomics* 4, 3084–3106.
- Pollack, J.H., Hashimoto, T., 1987. The role of glucose in the pH regulation of germ-tube formation in *Candida albicans*. *Journal of General Microbiology* 133, 415–424.
- Ramage, G., Saville, S.P., Wickes, B.L., López-Ribot, J.L., 2002. Inhibition of *Candida albicans* biofilm formation by farnesol, a quorum-sensing molecule. *Applied and Environmental Microbiology* 68, 5459–5463.
- Ramírez, M.A., Lorenz, M.C., 2007. Mutations in alternative carbon utilization pathways in *Candida albicans* attenuate virulence and confer pleiotropic phenotypes. *Eukaryotic Cell* 6, 280–290.
- Ramon, A.M., Porta, A., Fonzi, W.A., 1999. Effect of environmental pH on morphological development of *Candida albicans* is mediated via the PacC-related transcription factor encoded by PRR2. *Journal of Bacteriology* 181, 7524–7530.
- Rao, G.R., Sirsi, M., Ramakrishnan, T., 1962. Enzymes in *Candida albicans*. II. Tricarboxylic acid cycle and related enzymes. *Journal of Bacteriology* 84, 778–783.
- Reynolds, R., Braude, A.I., 1956. The filament inducing property of blood for *Candida albicans*; its nature and significance. *Clinical Research Procedure* 4, 40.
- Rocha, C.R.C., Schröppel, K., Marcus, D., Marcil, A., Dignard, D., Taylor, B.N., Thomas, D.Y., Whiteway, M., Leberer, E., 2001. Signaling through adenyl cyclase is essential for hyphal growth and virulence in the pathogenic fungus *Candida albicans*. *Molecular Biology of the Cell* 12, 3631–3643.
- Rodaki, A., Bohovych, I.M., Enjalbert, B., Young, T., Odds, F.C., Gow, N.A.R., Brown, A.J.P., 2009. Glucose promotes stress resistance in the fungal pathogen *Candida albicans*. *Molecular Biology of the Cell* 20, 4845–4855.
- Rogers, P.D., Barker, K.S., 2002. Evaluation of differential gene expression in fluconazole-susceptible and -resistant isolates of *Candida albicans* by cDNA microarray analysis. *Antimicrobial Agents and Chemotherapy* 46, 3412–3417.
- Román, E., Nombela, C., Pla, J., 2005. The Sho1 adaptor protein links oxidative stress to morphogenesis and cell wall biosynthesis in the fungal pathogen *Candida albicans*. *Molecular and Cellular Biology* 25, 10611–10627.
- Román, E., Arana, D.M., Nombela, C., Alonso-Monge, R., Pla, J., 2007. MAP kinase pathways as regulators of fungal virulence. *Trends in Microbiology* 15, 181–190.
- Román, E., Alonso-Monge, R., Gong, Q., Li, D., Calderone, R., Pla, J., 2009. The Cek1 MAPK is a short-lived protein regulated by quorum sensing in the fungal pathogen *Candida albicans*. *FEMS Yeast Research* 9, 942–955.
- Rooney, P.J., Klein, B.S., 2002. Linking fungal morphogenesis with virulence. *Cellular Microbiology* 4, 127–137.
- Rosenberg, E., Keller, K.H., Dworkin, M., 1977. Cell density dependent growth of *Myxococcus xanthus* on casein. *Journal of Bacteriology* 129, 770–777.
- Rubin-Bejerano, I., Fraser, I., Grisafi, P., Fink, G.R., 2003. Phagocytosis by neutrophils induces an amino acid deprivation response in *Saccharomyces cerevisiae* and *Candida albicans*. *Proceedings of the National Academy of Sciences of the United States of America* 100, 11007–11012.
- Sabie, F.T., Gadd, G.M., 1989. Involvement of a Ca^{2+} -calmodulin interaction in the yeast-mycelial (Y-M) transition of *Candida albicans*. *Mycopathologia* 108, 47–54.
- Sánchez-Martínez, C., Pérez-Martín, J., 2002. Gpa2, a G-protein subunit required for hyphal development in *Candida albicans*. *Eukaryotic Cell* 1, 865–874.
- Sato, T., Watanabe, T., Mikami, T., Matsumoto, T., 2004. Farnesol, a morphogenetic autoregulatory substance in the dimorphic fungus *Candida albicans*, inhibits hyphae growth through suppression of a mitogen-activated protein kinase cascade. *Biological and Pharmaceutical Bulletin* 27, 751–752.
- Saville, S.P., Lazzell, A.L., Monteagudo, C., Lopez-Ribot, J.L., 2003. Engineered control of cell morphology in vivo reveals distinct roles for yeast and filamentous forms of *Candida albicans* during infection. *Eukaryotic Cell* 2, 1053–1060.
- Schaefer, A.L., Val, D.L., Hanzelka, B.L., Cronan Jr., J.E., Greenberg, E.P., 1996. Generation of cell-to-cell signals in quorum sensing: acyl homoserine lactone synthase activity of a purified *Vibrio fischeri* LuxI protein. *Proceedings of the National Academy of Sciences of the United States of America* 93, 9505–9509.
- Scheper, M.A., Shirliff, M.E., Meiller, T.F., Peters, B.M., Jabra-Rizk, M.A., 2008. Farnesol, a fungal quorum-sensing molecule triggers apoptosis in human oral squamous carcinoma cells. *Neoplasia* 10, 954–963.
- Schwartz, D.S., Larsh, H.W., 1982. Comparative activities of glycolytic enzymes in yeast and mycelial forms of *Candida albicans*. *Mycopathologia* 78, 93–98.
- Schweizer, A., Rupp, S., Taylor, B.N., Rölinghoff, M., Schröppel, K., 2000. The TEA/ATTS transcription factor CaTc1p regulates hyphal development and virulence in *Candida albicans*. *Molecular Microbiology* 38, 435–445.
- Semighini, C.P., Hornby, J.M., Dumitru, R., Nickerson, K.W., Harris, S.D., 2006. Farnesol-induced apoptosis in *Aspergillus nidulans* reveals a possible mechanism for antagonistic interactions between fungi. *Molecular Microbiology* 59, 753–764.
- Sentheshanmuganathan, S., 1960. The mechanism of the formation of higher alcohols from amino acids by *Saccharomyces cerevisiae*. *Biochemical Journal* 47, 558–576.
- Sexton, J.A., Brown, V., Johnston, M., 2007. Regulation of sugar transport and metabolism by the *Candida albicans* Rgt1 transcriptional repressor. *Yeast* 24, 847–860.
- Shank, E.A., Kolter, R., 2009. New developments in microbial interspecies signaling. *Current Opinion in Microbiology* 12, 205–214.
- Sharkey, L.L., McNemar, M.D., Saporito-Irwin, S.M., Sypherd, P.S., Fonzi, W.A., 1999. HWP1 functions in the morphological development of *Candida albicans* downstream of EFG1, TUP1, and RBF1. *Journal of Bacteriology* 181, 5273–5279.
- Shchepin, R., Hornby, J.M., Burger, E., Niessen, T., Dussault, P., Nickerson, K.W., 2003. Quorum sensing in *Candida albicans*: probing farnesol's mode of action with 40 natural and synthetic farnesol analogs. *Chemistry and Biology* 10, 743–750.
- Shi, Q.M., Wang, Y.M., Zheng, X.D., Lee, R.T.H., Wang, Y., 2007. Critical role of DNA checkpoints in mediating genotoxic-stress-induced filamentous growth in *Candida albicans*. *Molecular Biology of the Cell* 18, 815–826.
- Shirliff, M.E., Krom, B.P., Meijering, R.A.M., Peters, B.M., Zhu, J., Scheper, M.A., Harris, M.L., Jabra-Rizk, M.A., 2009. Farnesol-induced apoptosis in *Candida albicans*. *Antimicrobial Agents and Chemotherapy* 53, 2392–2401.
- Sims, W., 1986. Effect of carbon dioxide on the growth and form of *Candida albicans*. *Journal of Medical Microbiology* 22, 203–208.
- Singh, V., Sinha, I., Sadhale, P.P., 2005. Global analysis of altered gene expression during morphogenesis of *Candida albicans* in vitro. *Biochemical and Biophysical Research Communications* 334, 1149–1158.
- Singh, V., Satheesh, S.V., Raghavendra, M.L., Sadhale, P.P., 2007. The key enzyme in galactose metabolism, UDP-galactose-4-epimerase, affects cell-wall integrity and morphology in *Candida albicans* even in the absence of galactose. *Fungal Genetics and Biology* 44, 563–574.
- Sinha, I., Wang, Y.M., Philp, R., Li, C.R., Yap, W.H., Wang, Y., 2007. Cyclin-dependent kinases control septin phosphorylation in *Candida albicans* hyphal development. *Developmental Cell* 13, 421–432.
- Smail, E.H., Cronstein, B.N., Meshulam, T., Esposito, A.L., Ruggeri, R.W., Diamond, R.D., 1992. In vitro, *Candida albicans* releases the immune modulator adenosine and a second, high-molecular weight agent that blocks neutrophil killing. *Journal of Immunology* 148, 3588–3595.
- Smith, D.A., Nicholls, S., Morgan, B.A., Brown, A.J.P., Quinn, J., 2004. A conserved stress-activated protein kinase regulates a core stress response in the human pathogen *Candida albicans*. *Molecular Biology of the Cell* 15, 4179–4190.
- Stoldt, V.R., Sonneborn, A., Leuker, C.E., Ernst, J.F., 1997. Efg1p, an essential regulator of morphogenesis of the human pathogen *Candida albicans*, is a member of a conserved class of bHLH proteins regulating morphogenetic processes in fungi. *EMBO Journal* 16, 1982–1991.
- Strich, R., Surosky, R.T., Steber, C., Dubois, E., Messenguy, F., Esposito, R.E., 1994. UME6 is a key regulator of nitrogen repression and meiotic development. *Genes and Development* 8, 796–810.
- Strijbis, K., Van Roermund, C.W.T., Hardy, G.P., Van Den Burg, J., Bloem, K., De Haan, J., Van Vlies, N., Wanders, R.J.A., Vaz, F.M., Distel, B., 2009. Identification and characterization of a complete carnitine biosynthesis pathway in *Candida albicans*. *FASEB Journal* 23, 2349–2359.
- Su, Z., Osborne, M.J., Xu, P., Xu, X., Li, Y., Ni, F., 2005. A bivalent dissection analysis of the high-affinity interactions between Cdc42 and the Cdc42/Rac interactive binding domains of signaling kinases in *Candida albicans*. *Biochemistry* 44, 16461–16474.
- Sudbery, P., Gow, N., Berman, J., 2004. The distinct morphogenic states of *Candida albicans*. *Trends in Microbiology* 12, 317–324.

- Sullivan, P.A., Shepherd, M.G., 1982. Gratuitous induction by N-acetylmannosamine of germ tube formation and enzymes for N-acetylglucosamine utilization in *Candida albicans*. *Journal of Bacteriology* 151, 1118–1122.
- Sun, J., Zeng, A.P., 2004. IdentiCS – identification of coding sequence and in silico reconstruction of the metabolic network directly from unannotated low-coverage bacterial genome sequence. *BMC Bioinformatics* 5.
- Swindell, K., Lattif, A.A., Chandra, J., Mukherjee, P.K., Ghannoum, M.A., 2009. Parenteral lipid emulsion induces germination of *Candida albicans* and increases biofilm formation on medical catheter surfaces. *Journal of Infectious Diseases* 200, 473–480.
- Swoboda, R.K., Bertram, G., Delbrück, S., Ernst, J.F., Gow, N.A.R., Gooday, G.W., Brown, A.J.P., 1994. Fluctuations in glycolytic mRNA levels during morphogenesis in *Candida albicans* reflect underlying changes in growth and are not a response to cellular dimorphism. *Molecular Microbiology* 13, 663–672.
- Tarry, W., Fisher, M., Shen, S., Mawhinney, M., 2005. *Candida albicans*: the estrogen target for vaginal colonization. *Journal of Surgical Research* 129, 278–282.
- Thomas, D.P., Viudes, A., Monteagudo, C., Lazzell, A.L., Saville, S.P., López-Ribot, J.L., 2006. A proteomic-based approach for the identification of *Candida albicans* protein components present in a subunit vaccine that protects against disseminated candidiasis. *Proteomics* 6, 6033–6041.
- Timpel, C., Strahl-Bolsinger, S., Ziegelbauer, K., Ernst, J.F., 1998. Multiple functions of Pmt1p-mediated protein O-mannosylation in the fungal pathogen *Candida albicans*. *Journal of Biological Chemistry* 273, 20837–20846.
- Timpel, C., Zink, S., Strahl-Bolsinger, S., Schröppel, K., Ernst, J., 2000. Morphogenesis, adhesive properties, and antifungal resistance depend on the Pmt6 protein mannosyltransferase in the fungal pathogen *Candida albicans*. *Journal of Bacteriology* 182, 3063–3071.
- Toenjes, K.A., Munsee, S.M., Ibrahim, A.S., Jeffrey, R., Edwards Jr., J.E., Johnson, D.I., 2005. Small-molecule inhibitors of the budded-to-hyphal-form transition in the pathogenic yeast *Candida albicans*. *Antimicrobial Agents and Chemotherapy* 49, 963–972.
- Tournu, H., Tripathi, G., Bertram, G., Macaskill, S., Mavor, A., Walker, L., Odds, F.C., Gow, N.A.R., Brown, A.J.P., 2005. Global role of the protein kinase Gcn2 in the human pathogen *Candida albicans*. *Eukaryotic Cell* 4, 1687–1696.
- Tripathi, G., Wiltshire, C., Macaskill, S., Tournu, H., Budge, S., Brown, A.J.P., 2002. Gcn4 co-ordinates morphogenetic and metabolic responses to amino acid starvation in *Candida albicans*. *EMBO Journal* 21, 5448–5456.
- Trunk, K., Gendron, P., Nantel, A., Lemieux, S., Roemer, T., Raymond, M., 2009. Depletion of the cullin Cdc53p induces morphogenetic changes in *Candida albicans*. *Eukaryotic Cell* 8, 756–767.
- Tsao, C.C., Chen, Y.T., Lan, C.Y., 2009. A small G protein Rhb1 and a GTPase-activating protein Tsc2 involved in nitrogen starvation-induced morphogenesis and cell wall integrity of *Candida albicans*. *Fungal Genetics and Biology* 46, 126–136.
- Urban, C., Sohn, K., Lottspeich, F., Brunner, H., Rupp, S., 2003. Identification of cell surface determinants in *Candida albicans* reveals Tsa1p, a protein differentially localized in the cell. *FEBS Letters* 544, 228–235.
- Vellucci, V.F., Gygas, S.E., Hostetter, M.K., 2007. Involvement of *Candida albicans* pyruvate dehydrogenase complex protein X (Pdx1) in filamentation. *Fungal Genetics and Biology* 44, 979–990.
- Vidotto, V., Accattatis, G., Zhang, Q., Campanini, G., Aoki, S., 1996a. Glucose influence on germ tube production in *Candida albicans*. *Mycopathologia* 133, 143–147.
- Vidotto, V., Sinicco, A., Accattatis, G., Aoki, S., 1996b. Influence of fructose on *Candida albicans* germ tube production. *Mycopathologia* 135, 85–88.
- Villas-Bôas, S.G., Roessner, U., Hansen, M.A.E., Smedsfaard, J., Nielsen, J., 2007. *Metabolome Analysis: An Introduction*. Wiley-Interscience, New Jersey.
- Wang, Y., 2009. CDKs and the yeast-hyphal decision. *Current Opinion in Microbiology* 12, 1–6.
- Wang, A., Raniga, P.P., Lane, S., Lu, Y., Liu, H., 2009. Hyphal chain formation in *Candida albicans*: Cdc28-Hgc1 phosphorylation of Efg1 represses cell separation genes. *Molecular and Cellular Biology* 29, 4406–4416.
- Warena, A.J., Kauffman, S., Sherrill, T.P., Becker, J.M., Konopka, J.B., 2003. *Candida albicans* septin mutants are defective for invasive growth and virulence. *Infection and Immunity* 71, 4045–4051.
- Westwater, C., Balish, E., Schofield, D.A., 2005. *Candida albicans*-conditioned medium protects yeast cells from oxidative stress: a possible link between quorum sensing and oxidative stress resistance. *Eukaryotic Cell* 4, 1654–1661.
- White, S., Larsen, B., 1997. *Candida albicans* morphogenesis is influenced by estrogen. *Cellular and Molecular Life Sciences* 53, 744–749.
- Whiteway, M., Bachewich, C., 2007. Morphogenesis in *Candida albicans*. *Annual Review of Microbiology* 61, 529–553.
- Wingard, J.R., Merz, W.G., Saral, R., 1979. *Candida tropicalis*: a major pathogen in immunocompromised patients. *Annals of Internal Medicine* 91, 539–543.
- Wisplinghoff, H., Bischoff, T., Tallent, S.M., Seifert, H., Wenzel, R.P., Edmond, M.B., 2004. Erratum: nosocomial bloodstream infections in US hospitals: analysis of 24, 179 cases from a prospective nationwide surveillance study (Clinical Infectious Diseases (August 1, 2004) 39 (309–317)). *Clinical Infectious Diseases* 39, 1093.
- Xu, X.-L., Lee, R.T.H., Fang, H.-M., Wang, Y.-M., Li, R., Zou, H., Zhu, Y., Wang, Y., 2008. Bacterial peptidoglycan triggers *Candida albicans* hyphal growth by directly activating the adenyl cyclase Cyr1p. *Cell Host & Microbe* 4, 28–39.
- Yamada-Okabe, T., Mio, T., Ono, N., Kashima, Y., Matsui, M., Arisawa, M., Yamada-Okabe, H., 1999. Roles of three histidine kinase genes in hyphal development and virulence of the pathogenic fungus *Candida albicans*. *Journal of Bacteriology* 181, 7243–7247.
- Yin, Z., Stead, D., Selway, L., Walker, J., Riba-Garcia, I., McInerney, T., Gaskell, S., Oliver, S.G., Cash, P., Brown, A.J.P., 2004. Proteomic response to amino acid starvation in *Candida albicans* and *Saccharomyces cerevisiae*. *Proteomics* 4, 2425–2436.
- Zaragoza, O., Blazquez, M.A., Gancedo, C., 1998. Disruption of the *Candida albicans* TPS1 gene encoding trehalose-6-phosphate synthase impairs formation of hyphae and decreases infectivity. *Journal of Bacteriology* 180, 3809–3815.
- Zheng, X., Wang, Y., 2004. Hgc1, a novel hypha-specific G1 cyclin-related protein regulates *Candida albicans* hyphal morphogenesis. *EMBO Journal* 23, 1845–1856.
- Zhou, H., Lorenz, M.C., 2008. Carnitine acetyltransferases are required for growth on non-fermentable carbon sources but not for pathogenesis in *Candida albicans*. *Microbiology* 154, 500–509.
- Zibafar, E., Hashemi, S.J., Zaini, F., Zeraati, H., Rezaie, S., Kordbacheh, P., 2009. Inhibitory effect of farnesol on biofilm formation by *Candida tropicalis*. *Daru* 17, 19–23.