



ELSEVIER

Available online at www.sciencedirect.com

SciVerse ScienceDirect

Current Opinion in
Cell Biology

Remodeling the nuclear membrane during closed mitosis

Dan Zhang^{1,2} and Snezhana Oliferenko^{1,2}

The mitotic spindle assembly and chromosome segregation in eukaryotes must be coordinated with the nuclear envelope (NE) remodeling. In a so-called 'open' mitosis the envelope of the mother nucleus is dismantled allowing the cytoplasmic spindle microtubules to capture the chromosomes. Alternatively, cells undergoing 'closed' mitosis assemble the intranuclear spindle and divide the nucleus without ever losing the nucleocytoplasmic compartmentalization. Here we focus on the mechanisms underlying mitotic NE dynamics in unicellular eukaryotes undergoing a closed nuclear division, paying specific attention to the emerging roles of the lipid biosynthesis machinery in this process. We argue that lessons learned in these organisms may be generally relevant to understanding the NE remodeling and the evolution of mitotic mechanisms throughout the eukaryotic domain.

Addresses

¹ Temasek Life Sciences Laboratory, 1 Research Link, Singapore 117604, Singapore

² Department of Biological Sciences, National University of Singapore, Singapore

Corresponding author: Oliferenko, Snezhana (snejana@tll.org.sg)

Current Opinion in Cell Biology 2012, 25:xx–yy

This review comes from a themed issue on **Cell architecture**

Edited by **Anna Akhmanova** and **Tim Stearns**

0955-0674/\$ – see front matter, © 2012 Elsevier Ltd. All rights reserved.

<http://dx.doi.org/10.1016/j.ceb.2012.09.001>

Introduction

Invention of the nucleus segregated from the cytoplasm by a selectively permeable membrane barrier has been a fundamental step in the advent of eukaryotic life. This called for integration of the nuclear membrane dynamics within the cellular physiology framework and in particular, its coordination with inheritance of the genetic material. Eukaryotes segregate duplicated chromosomes during mitosis using the microtubule-based spindle apparatus. This poses a spatial quandary since microtubules are normally located in the cytoplasm, where they function in intracellular transport and polarity establishment. Thus, either the nuclear membrane must break down early in mitosis to allow microtubules to capture the chromosomes or cells have to assemble an intranuclear mitotic spindle by importing the microtubule subunits and microtubule-associated proteins into an intact nucleus. The modern eukaryotes use both strategies – known as the open and the

closed mitosis – and a host of variations in between. This raises the questions of specific functional adaptations required for different types of mitoses, possible physiological advantages of distinct division modes and the degree of plasticity in this integral cell biological mechanism.

The nuclear envelope (NE) is an elaboration of the endomembrane system that consists of two lipid bilayers called the outer (ONM) and the inner nuclear membrane (INM). The ONM is continuous with the endoplasmic reticulum (ER), suggesting a close relationship between the NE and the general ER dynamics. The INM is occupied by a network of proteins interacting with the chromatin and contributing to the nuclear structure and function. The two NE membranes are fused at the nuclear pores, the sites of nucleocytoplasmic exchange decorated with multisubunit nuclear pore complexes (NPCs) [1]. Unlike the complex NE dynamics during open mitosis that involves NPC and nuclear lamina disassembly, release of the nuclear proteins and dispersal of the NE membranes into the ER, followed by their reassembly into the daughter nuclei [1,2], the closed nuclear division is essentially, a study in membrane remodeling. The major steps of this process include insertion of the microtubule-organizing centers (MTOCs) into the NE, anaphase nuclear membrane expansion and the fission of the nuclear membrane to yield two independent daughter nuclei. In this review, we will discuss mostly the membrane events preceding daughter nuclei individuation.

Membrane remodeling at the spindle pole body-nuclear envelope insertion sites

The spindle pole bodies (SPBs) serve as the MTOCs for the mitotic spindle in many lower eukaryotes. The SPB is a morphologically stratified structure duplicating once in each cell cycle [3,4]. Its inner plaque faces the nuclear interior and nucleates and anchors the spindle microtubules, whereas the outer plaque projects into the cytoplasm and nucleates the astral microtubule arrays. The central core ensures structural integrity of the SPB and anchors it within the NE plane. In budding yeast, the mother SPB localizes at the NE throughout the cell cycle and the newly formed daughter structure is inserted into the NE following the conservative SPB duplication in the G1 phase of the cell cycle [3,5,6]. **The daughter SPB insertion is driven by a concerted action of the SPB–NE tethering Mps3-Mps2-Bbp1-Nbp1 protein network [7–11] and the transmembrane nucleoporin Ndc1 that, incidentally, also contributes to the NPC anchorage at the NE [12].** Similarly to the NPC biogenesis, the SPB insertion machinery appears to function through the

2 Cell architecture

membrane-bending and membrane-anchoring properties of its constituents [11] although the molecular details of their action remain unknown.

Conceivably, the fusion between the INM and ONM resulting in the formation of a SPB pore, or 'fenestra', could also be promoted by local alterations in the spontaneous membrane curvature, for instance, by modulating distribution of lipids with fusogenic properties within the lipid bilayers [13]. The fission yeast *Schizosaccharomyces pombe* (*S. pombe*) perhaps provides a naturally sensitized system to study the SPB pore formation. Unlike in budding yeast, the *S. pombe* SPBs briefly settle within the NE during mitosis but are extruded to the outer nuclear face throughout the rest of the cell cycle [14] (Figure 1).


The NE fenestration and SPB insertion are well coordinated and a tight nucleocytoplasmic barrier is maintained throughout mitosis [15,16]. In cells carrying mutations in gene *brr6* the SPBs fail to properly anchor within the NE and the presumptive fenestration site remains agape [17^{••}]. *Brr6* encodes a member of the transmembrane NE protein family restricted to the organisms assembling the NE-spanning MTOCs and undergoing a 'closed' nuclear division. The Brr6 protein also appears to promote nuclear membrane remodeling during SPB extrusion at a later stage of mitosis.

In budding yeast, Brr6 and a functionally related protein Apq12 have been implicated in the NPC insertion into the NE, possibly through maintaining cellular lipid

homeostasis and affecting the nuclear membrane flexibility [18,19[•],20], although genetic interactions data suggest that these proteins may also function during the assembly of a 'closed' spindle [21]. Interestingly, treatment with chemicals that increase membrane fluidity alleviates the abnormal NE dynamics in cells lacking Apq12 [18,19[•]] but severely aggravates the Brr6-related phenotypes [19[•]]. Moreover, Apq12 and Brr6 mutants exhibit opposite patterns of sensitivity to growth at low and high temperatures. Since temperature shifts are known to trigger profound changes in the membrane fluidity through alterations in the length and the saturation status of the membrane phospholipid fatty acyl chains [22,23], Brr6 and Apq12 could function by either effecting the changes in membrane properties or possibly, sensing them. The recruitment of Brr6 to the mitotic SPBs that correlates with the SPB insertion and extrusion cycle, could perhaps argue for a direct, localized role of Brr6 in membrane lipid remodeling.

The precise molecular functions of Brr6 and Apq12 remain presently unknown but in light of the above observations, it would be of interest to see if direct manipulations of the fatty acid chain length and saturation status could affect the NE pore formation, during either the SPB or the NPC insertion events. The steady state fatty acid chain length distribution depends on the activity of the acetyl-CoA carboxylase that generates malonyl-CoA, the two-carbon donor in the synthesis of long-chain and very long-chain fatty acids. In budding yeast, the specific function of the acetyl-CoA carboxylase,

Figure 1



SPB-associated NE remodeling during mitosis in *S. pombe*. ONM, outer nuclear membrane; INM, inner nuclear membrane; CDK, cell cycle-dependent kinase. The resident SPB protein Cut12 boosts activity of Polo and CDK kinases specifically at the mitotic SPBs through amplification of the positive feedback loop. High activity of mitotic kinases in turn could promote localized remodeling of the NE and allow the SPB insertion. The red arrows indicate the presumptive signal for the NE fenestration.

Acc1, in the synthesis of very long-chain fatty acids is required for the normal NE structure [24,25]. Provocatively, Acc1 inactivation leads to a marked increase in the distance between the ONM and INM and a failure to properly insert the nuclear pores [24], possibly owing to defects in NE membrane fusion or pore maintenance. The *S. pombe* mutants of acetyl-CoA carboxylase exhibit severe spindle malfunction and fail to form equally sized daughter nuclei [26]. Reexamining the acetyl-CoA carboxylase mutants for the SPB insertion related phenotypes and testing the mutants that specifically affect synthesis of very long-chain fatty acids could prove useful in dissecting the NE membrane-shaping pathways. Similarly, the fatty acid saturation status could be modulated through the activity of the fatty acid desaturases and it would be of interest to examine possible effects of these enzymes on the SPB–NE interaction. Synthetic genetic interaction arrays and other currently available high-throughput methodologies should afford a systematic analysis of the functions of various lipid biosynthesis pathways in mitotic NE remodeling.

Cell cycle control of the SPB–NE insertion in fission yeast

The failure to integrate the SPBs in the *S. pombe* Brr6 mutants appears to parallel the consequences of mutating the Ndc1 ortholog Cut11 [27] and the mitotic regulator Cut12 [16,28]. While the effect of Cut11 deficiency is probably owing to a physical failure in anchoring the SPB within a fenestra, similar to its function in the NPC insertion, the reasons for the Cut12-related phenotype are less clear. Cut12 is an essential protein that lacks the membrane association domains; instead, it localizes to the inner edge of the SPB, facing the NE [28]. In cells carrying conditional loss-of-function alleles of *cut12*, the NE breaks down in the vicinity of the old, mother SPB but remains intact underneath the daughter SPB born in the previous interphase [16]. Thus, the old and the new SPBs apparently exhibit inherent differential sensitivity to the defects in the Cut12 function. Curiously, in spores germinating in the absence of Cut12, both SPBs fail to insert themselves into the NE or nucleate spindle microtubules [28].

A major biochemical role of Cut12 appears to promote mitosis through boosting the mitotic cyclin-dependent kinase (CDK) activity by amplification of the Polo kinase-driven positive feedback loop [29]. Both Polo [30,31] and the CDK complex [32] associate with the mitotic SPBs suggesting that the local activation of these kinases (through either a series or a parallel circuit) could promote localized NE fenestration and SPB insertion. Arguing for the involvement of the Polo kinase, conditional mutation in the fission yeast pericentrin Pcp1 leads to delocalization of Polo from the SPBs and a failure to fenestrate the NE in spite of an otherwise normal mitotic progression [33]. Furthermore, fenestra closure and the

SPB relocation to the outer nuclear surface at the end of anaphase coincide in time with the disappearance of Polo from the SPBs [30,31]. Evocative of the NE disassembly in higher eukaryotes, one could consider the SPB insertion in fission yeast essentially a localized NE breakdown triggered by activity of the mitotic kinases (Figure 1). Although the exact mechanisms relaying cell cycle signals to the SPB insertion remain to be elucidated, we predict that these may involve highly localized modulation of membrane properties leading to stabilization of the NE fenestra and the SPB anchorage by integral membrane proteins.


Scaling considerations in closed nuclear division

To accommodate spindle elongation and a closed division of the spherical mother nucleus into two daughters, the nuclear surface area increases during anaphase and it does so in a non-scalable manner [34]. For instance, a dividing *S. pombe* nucleus increases its surface area by approximately 30% by the end of mitosis while the nuclear volume remains constant [34,35]. This is not a simple issue of passively drawing the membrane material from the peripheral ER by the elongating nucleus. Arguing for a mitosis-specific mechanism of the NE growth, both *S. cerevisiae* and *S. pombe* cells deficient in spindle function and delaying in metaphase undergo extensive NE proliferation [36,37]. In cells with very little peripheral ER, such as a flagellate protist *Euglena*, the NE massively expands in preparation for mitosis, forming protuberances and invaginations that eventually resolve during nuclear division and formation of two spherical daughter nuclei [38].

The lipin family proteins that control the endomembrane growth appear to function at the crux of the matter (Figure 2). Interphase yeast cells lacking the lipin proteins (called Pah1 in *S. cerevisiae* and Ned1 in *S. pombe*) overproliferate both nuclear and peripheral ER membranes [39,40]. Enzymatically, lipins are Mg²⁺-dependent phosphatidate phosphatases that hydrolyze phosphatidic acid (PA) to diacylglycerol (DAG) [41]. Since triglycerides (TAG), the neutral storage lipids, are synthesized from DAG, the lack of Pah1 leads to a marked decrease in the cellular TAG levels [42]. In the absence of Pah1 function, PA can be rerouted through the CDP-DAG synthase pathway to yield the CDP-DAG, a key metabolite in the phospholipid production. In addition to its enzymatic function, Pah1 is involved in transcriptional repression of genes encoding the key phospholipid biosynthesis enzymes [39,43,44]. As a result, cells lacking Pah1 show a pronounced increase in steady state levels of structural phospholipids [42]. Modulating Pah1 activity can therefore balance the production of structural vs. storage lipids in many physiological scenarios, including the cell cycle progression. Indeed, Pah1 activity is subject to inhibitory phosphorylation by the cyclin-dependent

4 Cell architecture

Figure 2



Potential mechanistic links between lipin functions and the NE dynamics in yeast cells and the round worm, *C. elegans*. Solid lines indicate direct functions; dotted lines suggest indirect contributions. PA: phosphatidic acid; DAG: diacylglycerol.

kinases, including the mitotic CDK [45,46[•]]. The ER-localized Nem1/Spo7 phosphatase complex reverts the modification and promotes Pah1 activation [39,47,48]. Since the activity of CDK is highest at mitotic entry, the inhibition of Pah1 at this time point could account for a burst in phospholipid biosynthesis and assembly of a membrane reservoir. Curiously, lipins also function in NE remodeling during the open mitosis of the nematode *C. elegans* although the mechanisms underlying this phenomenon remain unclear [49–51].

Unlike the interphase cells lacking lipin function, the mitotic yeast cells do not overproliferate the entire ER network [36^{••}]. The preferential growth of the NE suggests the existence of a mechanism for mitosis-specific regulation of the lipid flow within the endomembrane system. One possibility could be a relatively localized assembly of the ER membranes competent for incorporation into the dividing nucleus. At least in case of *S. pombe*, cells appear to assemble the ER compartment enriched in proteins involved in membrane shaping and lipid metabolism at the equatorial region apposing the dividing nucleus ([52] and our unpublished data). It is also possible that the mitotic stage of the cell cycle is favorable for the growth of large ER sheets constituting the NE rather than the highly tubulated peripheral ER. The Ran GTPase system [15] and the NIMA kinase [53] have been implicated in regulating the NE integrity and/or expansion but the exact mechanisms underlying their functions remain unknown.


A pertinent question is where exactly the new membranes are inserted during the nuclear division. Studies in budding yeast suggest that the nuclear membrane adjacent to the nucleolar, chromatin-free region serves as an intrinsically expandable NE domain during non-scalable surface area increase [54]. The reasoning is that

the massive influx of new membranes throughout the NE could disrupt the complex organization of the chromosomal domains at the INM and adding a membrane ‘sink’ in a chromatin-free area could solve such problems [36^{••}]. However, the precise location of the ‘membrane injection’ sites in dividing yeast nuclei still awaits direct experimental characterization.

Striking diversity of mitotic mechanisms

The importance of assembling a membrane reservoir to enable the closed nuclear division is underscored by the unusual mitotic strategy of the fission yeast *Schizosaccharomyces japonicus* (*S. japonicus*) [35^{••},55[•]]. *S. japonicus* builds the intranuclear spindle and initiates the closed type of mitosis in a manner similar to its relative *S. pombe*. However, the *S. japonicus* nucleus fails to divide through the typical dumbbell intermediate. Instead, it elongates into diamond-like and bow-like shapes and spectacularly ruptures in late anaphase (Figure 3). The growing spindles initially buckle inside the intact nuclei but straighten upon the NE breakdown, suggesting that they can be constrained by the NE. Notably, the NE surface area in *S. japonicus* does not increase, and consistent with the simple scaling argument, the combined volume of the two re-formed daughter nuclei drops to approximately 70% of the mother [35^{••}]. NE rupture in *S. japonicus* is driven by the cell cycle machinery rather than by a mechanical force produced by the elongating spindle. When the membrane reservoir is made unavailable in *S. pombe* cells that do not have an intrinsic mechanism for the NE breakdown, the spindle buckles and breaks under the compressive stress and the diamond shaped mitotic nucleus collapses back to a single sphere of the original diameter [35^{••}]. Thus, it appears that at least within the fission yeast clade, the NE breakdown could have evolved to allow anaphase chromosome segregation and formation of two daughter nuclei in the absence of the mitotic NE growth. Whether

Figure 3



Nuclear envelope dynamics during mitotic progression in fission yeasts. A cartoon depicting the differences in the mitotic NE dynamics in *S. pombe* and *S. japonicus*.

such scaling considerations apply to evolution of mitotic mechanisms in all eukaryotes remains an open question.

The striking difference between the two related fission yeast species argues that invention of the variant mitotic mechanisms may require rather limited tinkering with the available cell physiology toolbox. This view is consistent with the unusual switch in mitotic NE behavior during development of *Physarum polycephalum*. During the amoeba stage, this slime mold assembles the centriole-based spindle and undergoes a typical open mitosis. However, upon commitment to the multinucleated plasmodium it undergoes a gradual transition to a fully closed mitosis [56].

What could be the physiological significance to the different types of NE behavior? Closed mitosis comes at the cost of producing a massive membrane reservoir in preparation for nuclear division and could possibly slow down the rates of spindle assembly and kinetochore capture in larger cells, owing to the necessity to import the entire microtubule machinery inside the intact nuclei. However, keeping the NE intact could prevent spindle entanglement, chromosome mis-segregation and uncontrolled nuclear fusion in a context of a multinucleated cell. This argument would fit well with the persistent nuclear membrane during mitosis in many filamentous ascomycetes and the early embryonic divisions in worms and flies that occur in syncytia. The dumbbell morphology of the dividing closed nucleus has been proposed to facilitate asymmetric segregation of nucleoplasmic proteins [57^{••}]

and circular DNA episomes [58^{••}] possibly contributing to functional differentiation within colonies of unicellular organisms. Finally, keeping the mitotic spindle – that helps to position the future cell division site in metazoans – within the enclosed compartment may provide a means for uncoupling chromosome segregation from cellular division plane specification.

Understanding the origins and functions of mitotic variation requires extending mechanistic analyses to multiple experimental systems, beyond the conventional few. A wealth of older ultrastructural records [59], combined with burgeoning genome sequence data, comprehensive phylogenetic analyses and the relative ease of experimentation, makes fungi especially attractive for this purpose. Establishing the rules and pathways for generating phenotypic diversity should provide unprecedented insights into the fundamental aspects of nuclear evolution.

Acknowledgements

We are grateful to E. Makeyev and A. Vjestica for discussions and suggestions on the manuscript. Our work is supported by Singapore Millenium Foundation.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Hetzer MW: **The nuclear envelope**. *Cold Spring Harb Perspect Biol* 2010, **2**:a000539.

6 Cell architecture

2. Guttinger S, Laurell E, Kutay U: **Orchestrating nuclear envelope disassembly and reassembly during mitosis.** *Nat Rev Mol Cell Biol* 2009, **10**:178-191.
3. O'Toole ET, Winey M, McIntosh JR: **High-voltage electron tomography of spindle pole bodies and early mitotic spindles in the yeast *Saccharomyces cerevisiae*.** *Mol Biol Cell* 1999, **10**:2017-2031.
4. Bullitt E, Rout MP, Kilmartin JV, Akey CW: **The yeast spindle pole body is assembled around a central crystal of Spc42p.** *Cell* 1997, **89**:1077-1086.
5. Byers B, Goetsch L: **Behavior of spindles and spindle plaques in the cell cycle and conjugation of *Saccharomyces cerevisiae*.** *J Bacteriol* 1975, **124**:511-523.
6. Adams IR, Kilmartin JV: **Localization of core spindle pole body (SPB) components during SPB duplication in *Saccharomyces cerevisiae*.** *J Cell Biol* 1999, **145**:809-823.
7. Schramm C, Elliott S, Shevchenko A, Schiebel E: **The Bbp1p-Mps2p complex connects the SPB to the nuclear envelope and is essential for SPB duplication.** *EMBO J* 2000, **19**:421-433.
8. Araki Y, Lau CK, Maekawa H, Jaspersen SL, Giddings TH, Schiebel E, Winey M: **The *Saccharomyces cerevisiae* spindle pole body (SPB) component Nbp1p is required for SPB membrane insertion and interacts with the integral membrane proteins Ndc1p and Mps2p.** *Mol Biol Cell* 2006, **17**:1959-1970.
9. Kupke T, Di Cecco L, Muller HM, Neuner A, Adolf F, Wieland F, Nickel W, Schiebel E: **Targeting of Nbp1 to the inner nuclear membrane is essential for spindle pole body duplication.** *EMBO J* 2011, **30**:3337-3352.
10. Friederichs JM, Ghosh S, Smoyer CJ, McCroskey S, Miller BD, Weaver KJ, Delventhal KM, Unruh J, Slaughter BD, Jaspersen SL: **The SUN protein Mps3 is required for spindle pole body insertion into the nuclear membrane and nuclear envelope homeostasis.** *PLoS Genet* 2011, **7**:e1002365.
11. Jaspersen SL, Ghosh S: **Nuclear envelope insertion of spindle pole bodies and nuclear pore complexes.** *Nucleus* 2012, **3**:226-236.
12. Lau CK, Giddings TH, Winey M: **A novel allele of *Saccharomyces cerevisiae* NDC1 reveals a potential role for the spindle pole body component Ndc1p in nuclear pore assembly.** *Eukaryot Cell* 2004, **3**:447-458.
13. Chernomordik LV, Kozlov MM: **Mechanics of membrane fusion.** *Nat Struct Mol Biol* 2008, **15**:675-683.
14. Ding R, West RR, Morphew DM, Oakley BR, McIntosh JR: **The spindle pole body of *Schizosaccharomyces pombe* enters and leaves the nuclear envelope as the cell cycle proceeds.** *Mol Biol Cell* 1997, **8**:1461-1479.
15. Gonzalez Y, Meerbrey K, Chong J, Torii Y, Padte NN, Sazer S: **Nuclear shape, growth and integrity in the closed mitosis of fission yeast depend on the Ran-GTPase system, the spindle pole body and the endoplasmic reticulum.** *J Cell Sci* 2009, **122**:2464-2472.
16. Tallada VA, Tanaka K, Yanagida M, Hagan IM: **The *S. pombe* mitotic regulator Cut12 promotes spindle pole body activation and integration into the nuclear envelope.** *J Cell Biol* 2009, **185**:875-888.
17. Tamm T, Grallert A, Grossman EP, Alvarez-Tabares I, Stevens FE, Hagan IM: **Brr6 drives the *Schizosaccharomyces pombe* spindle pole body nuclear envelope insertion/extrusion cycle.** *J Cell Biol* 2011, **195**:467-484.
- This interesting paper demonstrates that the NE protein Brr6 previously shown to function in maintaining nuclear membrane fluidity and homeostasis is required for the NE insertion and extrusion of the mitotic SPBs in *S. pombe*.
18. Scarcelli JJ, Hodge CA, Cole CN: **The yeast integral membrane protein Apq12 potentially links membrane dynamics to assembly of nuclear pore complexes.** *J Cell Biol* 2007, **178**:799-812.
19. Hodge CA, Choudhary V, Wolyniak MJ, Scarcelli JJ, Schneider R, Cole CN: **Integral membrane proteins Brr6 and Apq12 link assembly of the nuclear pore complex to lipid homeostasis in the endoplasmic reticulum.** *J Cell Sci* 2010, **123**:141-151.
- This study suggests how the NE integral membrane proteins Brr6 and Apq12 could function in nuclear pore complex assembly and anchorage in *S. cerevisiae*.
20. Schneider R, Cole CN: **Integrating complex functions: coordination of nuclear pore complex assembly and membrane expansion of the nuclear envelope requires a family of integral membrane proteins.** *Nucleus* 2010, **1**:387-392.
21. Montpetit B, Thorne K, Barrett I, Andrews K, Jadusingsh R, Hieter P, Measday V: **Genome-wide synthetic lethal screens identify an interaction between the nuclear envelope protein, Apq12p, and the kinetochore in *Saccharomyces cerevisiae*.** *Genetics* 2005, **171**:489-501.
22. Los DA, Murata N: **Membrane fluidity and its roles in the perception of environmental signals.** *Biochim Biophys Acta* 2004, **1666**:142-157.
23. Zhang YM, Rock CO: **Membrane lipid homeostasis in bacteria.** *Nat Rev Microbiol* 2008, **6**:222-233.
24. Schneider R, Hitomi M, Ivessa AS, Fasch EV, Kohlwein SD, Tartakoff AM: **A yeast acetyl coenzyme A carboxylase mutant links very-long-chain fatty acid synthesis to the structure and function of the nuclear membrane-pore complex.** *Mol Cell Biol* 1996, **16**:7161-7172.
25. Al-Feel W, DeMar JC, Wakil SJ: **A *Saccharomyces cerevisiae* mutant strain defective in acetyl-CoA carboxylase arrests at the G2/M phase of the cell cycle.** *Proc Natl Acad Sci USA* 2003, **100**:3095-3100.
26. Saitoh S, Takahashi K, Nabeshima K, Yamashita Y, Nakaseko Y, Hirata A, Yanagida M: **Aberrant mitosis in fission yeast mutants defective in fatty acid synthetase and acetyl CoA carboxylase.** *J Cell Biol* 1996, **134**:949-961.
27. West RR, Vaisberg EV, Ding R, Nurse P, McIntosh JR: **Cut11(+): a gene required for cell cycle-dependent spindle pole body anchoring in the nuclear envelope and bipolar spindle formation in *Schizosaccharomyces pombe*.** *Mol Biol Cell* 1998, **9**:2839-2855.
28. Bridge AJ, Morphew M, Bartlett R, Hagan IM: **The fission yeast SPB component Cut12 links bipolar spindle formation to mitotic control.** *Genes Dev* 1998, **12**:927-942.
29. Hagan IM: **The spindle pole body plays a key role in controlling mitotic commitment in the fission yeast *Schizosaccharomyces pombe*.** *Biochem Soc Trans* 2008, **36**:1097-1101.
30. Bahler J, Steever AB, Wheatley S, Wang Y, Pringle JR, Gould KL, McCollum D: **Role of polo kinase and Mid1p in determining the site of cell division in fission yeast.** *J Cell Biol* 1998, **143**:1603-1616.
31. Mulvihill DP, Petersen J, Ohkura H, Glover DM, Hagan IM: **Plo1 kinase recruitment to the spindle pole body and its role in cell division in *Schizosaccharomyces pombe*.** *Mol Biol Cell* 1999, **10**:2771-2785.
32. Decottignies A, Zarzov P, Nurse P: **In vivo localisation of fission yeast cyclin-dependent kinase cdc2p and cyclin B cdc13p during mitosis and meiosis.** *J Cell Sci* 2001, **114**:2627-2640.
33. Fong CS, Sato M, Toda T: **Fission yeast Pcp1 links polo kinase-mediated mitotic entry to gamma-tubulin-dependent spindle formation.** *EMBO J* 2010, **29**:120-130.
- The authors show that in *S. pombe*, the pericentriin protein Pcp1 functions in recruiting the Polo kinase to the mitotic SPBs and promoting the SPB insertion into the NE.
34. Lim HWG, Huber G, Torii Y, Hirata A, Miller J, Sazer S: **Vesicle-like biomechanics governs important aspects of nuclear geometry in fission yeast.** *PLoS ONE* 2007, **2**:e948.
35. Yam C, He Y, Zhang D, Chiam KH, Oliferenko S: **Divergent strategies for controlling the nuclear membrane satisfy geometric constraints during nuclear division.** *Curr Biol* 2011, **21**:1314-1319.
- This study suggests that control of the nuclear membrane area may determine the choice between the NE breakdown and the 'closed' nuclear division, at least within the fission yeast clade.

36. Witkin KL, Chong Y, Shao S, Webster MT, Lahiri S, Walters AD, ●● Lee B, Koh JL, Prinz WA, Andrews BJ *et al.*: **The budding yeast nuclear envelope adjacent to the nucleolus serves as a membrane sink during mitotic delay.** *Curr Biol* 2012, **22**:1128-1133.
This study highlights a massive and spatially delimited expansion of the nuclear membrane during mitotic delay in budding yeast.
37. Castagnetti S, Oliferenko S, Nurse P: **Fission yeast cells undergo nuclear division in the absence of spindle microtubules.** *PLoS Biol* 2010, **8**:e1000512.
38. Gillott MA, Triemer RE: **The ultrastructure of cell division in *Euglena gracilis*.** *J Cell Sci* 1978, **31**:25-35.
39. Santos-Rosa H, Leung J, Grimsey N, Peak-Chew S, Siniosoglou S: **The yeast lipin Smp2 couples phospholipid biosynthesis to nuclear membrane growth.** *EMBO J* 2005, **24**:1931-1941.
40. Tange Y, Hirata A, Niwa O: **An evolutionarily conserved fission yeast protein, Ned1, implicated in normal nuclear morphology and chromosome stability, interacts with Dis3, Pim1/RCC1 and an essential nucleoporin.** *J Cell Sci* 2002, **115**:4375-4385.
41. Han GS, Wu WJ, Carman GM: **The *Saccharomyces cerevisiae* Lipin homolog is a Mg²⁺-dependent phosphatidate phosphatase enzyme.** *J Biol Chem* 2006, **281**:9210-9218.
42. Carman GM, Han GS: **Roles of phosphatidate phosphatase enzymes in lipid metabolism.** *Trends Biochem Sci* 2006, **31**:694-699.
43. O'Hara L, Han GS, Peak-Chew S, Grimsey N, Carman GM, Siniosoglou S: **Control of phospholipid synthesis by phosphorylation of the yeast lipin Pah1p/Smp2p Mg²⁺-dependent phosphatidate phosphatase.** *J Biol Chem* 2006, **281**:34537-34548.
44. Loewen CJ, Gaspar ML, Jesch SA, Delon C, Ktistakis NT, Henry SA, Levine TP: **Phospholipid metabolism regulated by a transcription factor sensing phosphatidic acid.** *Science* 2004, **304**:1644-1647.
45. Choi HS, Su WM, Morgan JM, Han GS, Xu Z, Karanasios E, Siniosoglou S, Carman GM: **Phosphorylation of phosphatidate phosphatase regulates its membrane association and physiological functions in *Saccharomyces cerevisiae*: identification of SER(602), THR(723), AND SER(744) as the sites phosphorylated by CDC28 (CDK1)-encoded cyclin-dependent kinase.** *J Biol Chem* 2011, **286**:1486-1498.
46. Karanasios E, Han GS, Xu Z, Carman GM, Siniosoglou S: ●● **A phosphorylation-regulated amphipathic helix controls the membrane translocation and function of the yeast phosphatidate phosphatase.** *Proc Natl Acad Sci USA* 2010, **107**:17539-17544.
This study provides a mechanistic basis for understanding functions of lipin proteins in regulating NE and ER structure by showing how the budding yeast lipin Pah is activated and translocates to the ER membranes.
47. Siniosoglou S, Santos-Rosa H, Rappsilber J, Mann M, Hurt E: **A novel complex of membrane proteins required for formation of a spherical nucleus.** *EMBO J* 1998, **17**:6449-6464.
48. Kim Y, Gentry MS, Harris TE, Wiley SE, Lawrence JC, Dixon JE: **A conserved phosphatase cascade that regulates nuclear membrane biogenesis.** *Proc Natl Acad Sci USA* 2007, **104**:6596-6601.
49. Gorjanacz M, Mattaj IW: **Lipin is required for efficient breakdown of the nuclear envelope in *Caenorhabditis elegans*.** *J Cell Sci* 2009, **122**:1963-1969.
50. Golden A, Liu J, Cohen-Fix O: **Inactivation of the *C. elegans* lipin homolog leads to ER disorganization and to defects in the breakdown and reassembly of the nuclear envelope.** *J Cell Sci* 2009, **122**:1970-1978.
51. Han S, Bahmanyar S, Zhang P, Grishin N, Oegema K, Crooke R, Graham M, Reue K, Dixon JE, Goodman JM: **Nuclear envelope phosphatase 1-regulatory subunit 1 (formerly TMEM188) is the metazoan Spo7p ortholog and functions in the lipin activation pathway.** *J Biol Chem* 2012, **287**:3123-3137.
52. Zhang D, Vjestica A, Oliferenko S: **The cortical ER network limits the permissive zone for actomyosin ring assembly.** *Curr Biol* 2010, **20**:1029-1034.
53. Krien MJ, West RR, John UP, Koniaras K, McIntosh JR, O'Connell MJ: **The fission yeast NIMA kinase Fin1p is required for spindle function and nuclear envelope integrity.** *EMBO J* 2002, **21**:1713-1722.
54. Campbell JL, Lorenz A, Witkin KL, Hays T, Loidl J, Cohen-Fix O: **Yeast nuclear envelope subdomains with distinct abilities to resist membrane expansion.** *Mol Biol Cell* 2006, **17**:1768-1778.
55. Aoki K, Hayashi H, Furuya K, Sato M, Takagi T, Osumi M, Kimura A, ●● Niki H: **Breakage of the nuclear envelope by an extending mitotic nucleus occurs during anaphase in *Schizosaccharomyces japonicus*.** *Genes Cells* 2011, **16**:911-926.
This paper provides interesting ultrastructural and cell biological analyses of the variant mitosis in *S. japonicus*.
56. Burland TG, Solnica-Krezel L, Bailey J, Cunningham DB, Dove WF: **Patterns of inheritance, development and the mitotic cycle in the protist *Physarum polycephalum*.** *Adv Microb Physiol* 1993, **35**:1-69.
57. Boettcher B, Marquez-Lago TT, Bayer M, Weiss EL, Barral Y: ●● **Nuclear envelope morphology constrains diffusion and promotes asymmetric protein segregation in closed mitosis.** *J Cell Biol* 2012, **197**:921-937.
This paper suggests that the dumbbell-like morphology of the dividing nucleus in *S. cerevisiae* could promote asymmetric segregation of nucleoplasmic proteins.
58. Gehlen LR, Nagai S, Shimada K, Meister P, Taddei A, Gasser SM: ●● **Nuclear geometry and rapid mitosis ensure asymmetric episome segregation in yeast.** *Curr Biol* 2011, **21**:25-33.
The authors show that the characteristic shape of the dividing nucleus and the short duration of mitosis may restrict passive diffusion of plasmids into daughter nuclei in *S. cerevisiae*.
59. Heath IB: **Variant mitoses in lower eukaryotes: indicators of the evolution of mitosis.** *Int Rev Cytol* 1980, **64**:1-80.